

Positive Maternal Mental Health During Pregnancy Associated With Specific Forms of Adaptive Development in Early Childhood: Evidence from a Longitudinal Study

Desiree Y Phua¹, Michelle KL Kee¹, Dawn X.P. Koh¹, Anne Rifkin-Graboi¹, Mary Daniels², Helen Chen², Yap Seng Chong^{1,3}, Peter D Gluckman^{1,4}, Birit FP Broekman¹, Iliana Magiati⁵, Neerja Karnani^{1,6}, Michael Pluess⁷ and Michael J Meaney^{1,8} on behalf of the GUSTO study Group*

1. Singapore Institute for Clinical Sciences (SICS), A*STAR, Brenner Centre for Molecular Medicine, Singapore.
2. KK Women's and Children's Hospital, Singapore.
3. Department of Obstetrics and Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore (NUS), Singapore.
4. Centre for Human Evolution, Adaptation and Disease, Liggins Institute, University of Auckland, Auckland, New Zealand.
5. Department of Psychology, National University of Singapore, Singapore 117576, Singapore.
6. Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore (NUS), Singapore.
7. Department of Biological and experimental Psychology, Queen Mary University London, United Kingdom E1 4NS
8. Department of Psychiatry, Douglas Mental Health University Institute, McGill University, Montreal, Canada H4H 1R3
9. Sackler Program for Epigenetics & Psychobiology at McGill University, Montreal, Canada H4H 1R3

Correspondence: Michael J Meaney

Sackler Program for Epigenetics & Psychobiology at McGill University

Douglas Mental Health University Institute

6875 boul LaSalle

Montreal, Canada H4H 1R3

e.mail: michael.meaney@mcgill.ca

Acknowledgements

The GUSTO study is funded by the Singapore National Research Foundation under its Translational and Clinical Research (TCR) Flagship Programme and administered by the Singapore Ministry of Health's National Medical Research Council (NMRC), Singapore-NMRC/TCR/004-NUS/2008; NMRC/TCR/012-NUHS/2014. Additional funding is provided by the Singapore Institute for Clinical Sciences, Agency for Science Technology and Research (A*STAR), Singapore.. We acknowledge additional funding from the Sackler Foundation (MJM). We thank the GUSTO study group and all clinical and home visit staff involved. The voluntary participation of all participants is greatly appreciated.

*The GUSTO study group includes Pratibha Agarwal, Arijit Biswas, Choon Looi Bong, Shirong Cai, Jerry Kok Yen Chan, Yiong Huak Chan, Cornelia Yin Ing Chee, Yin Bun Cheung, Audrey Chia, Amutha Chinnadurai, Chai Kiat Chng, Mary Foong-Fong Chong, Shang Chee Chong, Mei Chien Chua, Chun Ming Ding, Eric Andrew Finkelstein, Doris Fok, Keith M. Godfrey, Anne Eng Neo Goh, Yam Thiam Daniel Goh, Joshua J. Gooley, Wee Meng Han, Mark Hanson, Christiani Jeyakumar Henry, Joanna D. Holbrook, Chin-Ying Hsu, Hazel Inskip, Jeevesh Kapur, Ivy Yee-Man Lau, Bee Wah Lee, Yung Seng Lee, Ngee Lek, Sok Bee Lim, Yen-Ling Low, Iliana Magiati, Lourdes Mary Daniel, Cheryl Ngo, Krishnamoorthy Naiduvaje, Wei Wei Pang, Boon Long Quah, Victor Samuel Rajadurai, Mary Rauff, Salome A. Rebello, Jenny L. Richmond, Lynette Pei-Chi Shek, Allan Sheppard, Borys Shuter, Leher Singh, Shu-E Soh,

Walter Stunkel, Lin Lin Su, Kok Hian Tan, Oon Hoe Teoh, Mya Thway Tint, Hugo P S van Bever, Rob M. van Dam, Inez Bik Yun Wong, P. C. Wong, Fabian Yap, George Seow Heong Yeo.

Abstract

The quality of prenatal maternal mental health, from psychological stress, depressive symptoms, to anxiety and other non-psychotic mental disorders profoundly affects fetal neurodevelopment. Despite the evidence for the influence of positive mental well-being on health there is, to our knowledge, no research examining the possible effects of positive antenatal mental health on the development of the offspring. Using exploratory bifactor analysis, this prospective study ($n = 1066$) demonstrated the feasibility of using common psychiatric screening tools to examine the effect of positive maternal mental health. Antenatal mental health was assessed during 26-week pregnancy. The effects on offspring was assessed when the child was 12-, 18- and 24-months-old. Results showed that positive antenatal mental health was uniquely associated with the offspring's cognitive, language and parentally-rated competences. This study shows that the effects of positive maternal mental health are likely to be specific and different from the lack of mental disorders.

Introduction

The quality of prenatal maternal mental health, from psychological stress (e.g., Beydoun & Saftlas, 2008; Charil, Laplante, Vaillancourt, & King, 2010; Graignic-Philippe, Dayan, Chokron, Jacquet, & Tordjman, 2014), depressive symptoms (e.g., Field, 2011; Gentile, 2017; Mulder et al., 2002; Waters, Hay, Simmonds, & van Goozen, 2014), to anxiety (e.g., Van den Bergh, Mulder, Mennes, & Glover, 2005) and other non-psychotic mental disorders (Howard et al., 2014) profoundly affects fetal neurodevelopment. Such effects are apparent in terms of neural structure and organization (Buss et al., 2012; Qiu, Tuan, Li, et al., 2015; Qiu, Tuan, Ong, et al., 2015; Rifkin-Graboi et al., 2015), cognitive and emotional function as well as the subsequent risk for psychopathology (Baibazarova et al., 2013; Goodman et al., 2011; Graignic-Philippe et al., 2014; O'Donnell & Meaney, 2017; Pluess et al., 2011). The effects of maternal conditions can even be transmitted to the third generation (e.g., Babenko, Kovalchuk, & Metz, 2015; Bowers & Yehuda, 2016; Gröger et al., 2016). Importantly, the effects of prenatal maternal mental health persist even after controlling for postnatal maternal status (Glover, 2014; Huizink, Mulder, & Buitelaar, 2004; Pearson et al., 2013). In the case of depression, the effects of prenatal maternal states appear to be statistically more strongly associated with the later risk of depression in the offspring that are those of postnatal maternal depressive symptoms (Pearson et al., 2013). A 'prenatal cross-fostering' study in humans where pregnant mothers were related or unrelated to their child as a result of in vitro fertilization, which served to distinguish maternally-inherited effects from those directly associated with the maternal phenotype, showed that maternal stress and emotional well-being were directly associated with socio-emotional function in the child (Rice et al., 2010).

While about 12-15% of pregnant women screen positively for depression (e.g., Bennett, Einarson, Taddio, Koren, & Einarson, 2004; Gavin et al., 2005; Karmaliani et al., 2009; Le Strat, Dubertret, & Le Foll, 2011), there is substantial variation in the psychological well-being among

the remaining mothers (Keyes, 2002). Neuroimaging studies, including those performed with neonates, show that the influence of symptoms of anxiety and depression cuts across the entire population and are not unique to the offspring of mothers with confirmed clinical disorders (Buss et al., 2012; Qiu, Tuan, Li, et al., 2015; Qiu, Tuan, Ong, et al., 2015; Rifkin-Graboi et al., 2015). The same finding emerges from studies of a wide range of neurodevelopmental outcomes. Despite the compelling evidence for the broad influence of maternal emotional well-being, the existing literature focuses almost exclusively on the effects of stress or symptoms of depression or anxiety, and does therefore not capture the full range of mental well-being. The potential effect of positive antenatal mental health on neurodevelopment in the offspring will allow us to examine the broader spectrum of mental health and consider promoting health rather than merely preventing mental disorders.

Positive mental health

Health is a continuum that includes a sense of well-being and is not merely defined by the absence of illness or disability (World Health Organization, 2004). Positive and negative mental health, though correlated, are distinct constructs (Huppert & Whittington, 2003). Effective interventions may reduce depressive symptoms, but do little to increase mental well-being (Newnham, Hooke, & Page, 2010), again suggesting the independence of both constructs (de Cates, Stranges, Blake, & Weich, 2015). Furthermore, positive mental health and mental illness symptoms have different antecedents including various demographics and socio-emotional variables (Hu, Stewart-Brown, Twigg, & Weich, 2007).

While positive antenatal mental health has been largely neglected, existing studies show that a higher level of mental well-being serves as a protective factor against future mental disorders (Keyes, Dhingra, & Simoes, 2010; Lamers, Westerhof, Glas, & Bohlmeijer, 2015). This protective factor is apparent in children; while paternal depression strongly predicted depressive symptoms, this effect was not seen in children with positive mental health traits (Tam

et al., 2017). Positive mental health in young adulthood can even predict a range of health outcomes (Aspinwall & Tedeschi, 2010; Howell, Kern, & Lyubomirsky, 2007) as well as mortality in late adulthood (Danner, Snowdon, & Friesen, 2001). Despite the evidence for the influence of positive mental well-being on health there is, to our knowledge, no research examining the possible effects of positive antenatal mental health on the development of the offspring.

Bi-factor model of maternal mental health symptoms

While large-scale birth-cohort studies emphasize the importance of maternal mental health problems, measures of positive mental health in the study design are rarely considered. The most commonly used measures of maternal mental health focus on symptoms of depression (e.g., Edinburgh Postnatal Depression Scale, Center for Epidemiologic Studies Depression Scale) or anxiety (e.g., State-Trait Anxiety Inventory, anxiety subscale of Crown Crisp Experiential Index). While such measures are used to screen for symptoms of mental disorders, it may nevertheless be possible to detect aspects of positive mental health. For example, though the General Health Questionnaire is a psychiatric disorder screening tool, Hu and colleagues (2007) used factor analyses to show that the positively-worded items can be indicators of positive mental health and not merely absence of symptoms of mental disorders. The STAI has likewise been used to reflect positive mental health or well-being as well (Hernández-Martínez, Val, Murphy, Busquets, & Sans, 2011; Kvaal, Laake, & Engedal, 2001). Thus psychiatric disorders screening tools appear to contain items that reflect positive mental health.

Bi-factor modelling is increasingly used to factor analyze the multidimensional nature of mental health. The premise of bi-factor modelling is that there is an overarching general mental health or psychopathology dimension or factor that reflects responses to the mental health measures regardless of the nature of disorder (e.g., Caspi et al., 2014; Simms, Grös, Watson, & O'hara, 2008). There is therefore considerable value to the inclusion of multiple measures of

mental health within a single bi-factor latent model. There is heterogeneity in antenatal mental health, even if only focused on antenatal depression (Castro et al., 2016; Santos, Tan, & Salomon, 2016). Mental health instruments are often checklists of symptoms that are commonly associated with specific disorders. However, even within measures of depression, there is substantial breadth to the features that are examined (Fried, 2017), which provide a more comprehensive analysis of maternal mental health. In this paper we report the results of bi-factor analysis using data from a longitudinal birth cohort with multiple, commonly-used measures of the symptoms of anxiety or depression in women at mid-gestation. The results yielded coherent measures of antenatal positive mental health that predicted developmental outcomes in the children, especially those focusing on social behaviors and communication.

Methods

Participants

This study was part of a prospective birth cohort study, Growing Up in Singapore Towards Healthy Outcomes (GUSTO; see Soh et al., 2012). The GUSTO sample ($n = 1066$) included women who conceived naturally (i.e., not through in-vitro fertilization), did not have any medical conditions before or during pregnancy, and gave birth to single full-term babies (i.e., non-twins) with normal birth weight. After delivery, the participants and their children were invited to the study clinic at when the child was 12 months, 18 months and 24 months old. At the clinic, the child was administered a battery of neurocognitive and behavioral tasks. The mothers were also given measures about their child's behavior.

Scales

Maternal mental health. Three maternal mental health measures were administered during 26-week pregnancy during the participants' regular clinic visit. The responses to the individual items of the measures were used in the bi-factor models.

The Beck Depression Inventory II (BDI-II; Beck, Steer, & Brown, 1996) is an inventory of 21 cluster of items describing common depressive symptoms. Each cluster contains four to seven statements describing varying severity of a common depressive symptom (e.g., feeling worthless). Participants selected the statement that best described how they felt for the past two weeks. The Edinburgh Postnatal Depression Scale (EPDS; Cox, Holden, & Sagovsky, 1987) has ten items of depressive symptoms and participants indicated how much each item described how they were feeling for past 7 days on a 4-point Likert scale. The State-Trait Anxiety Inventory (STAI; Spielberger, Gorsuch, & Lushene, 1970) consisted of 40 items that are associated with anxiety (or lack of). For the first 20 items, participants responded to how much each item described how they felt right now on a 4-point Likert scale; for the next 20 items, they responded to how much the item described how they generally felt.

Child measures. When the child was 12-month old, mothers rated their child's socio-emotional behavior on the Infant Toddler Socio-Emotional Assessment (ITESA) questionnaire (Briggs-Gowan & Carter, 1998). Twenty-one behaviors on four domains (internalizing, externalizing, dysregulation, & competence behaviors) were assessed (see Table 6 for the list of behaviors).

At 18-month of age, mothers rated their child's behavior on the 25-item Quantitative Checklist for Autism in Toddlers (QChat; Allison et al., 2008). Other than a total score, there were two subscores corresponding to the behavioral and social factors of autism-traits.

At 24-month, the Bayley Scales of Infant and Toddler Development (Bayley), third edition (Bayley, 2006) was used to assess the child's development in the domains of cognition, language, motor skills, socio-emotional behaviors and adaptability. The cognitive, language (i.e., receptive and expressive communication) and motor skills (i.e., fine and gross motor) components were assessed via standardized laboratory tasks. The socio-emotional and adaptability (i.e., Communication behavior, community use, functional pre-academic, home

living, health and safety knowledge, leisure activities, self-care, self-direction skills, social skills, and motor skills) were rated by their care-giver.

Statistical analyses

Bi-factor models. An exploratory bi-factor model was first fitted with the individual items of the mental health scales as manifest variables. Number of factors was determined with parallel analysis. In parallel analysis, eigenvalues from randomly generated correlation matrices were computed. A factor will be retained if the eigenvalue from the observed data is larger than the corresponding eigenvalue from parallel analysis (Hayton, Allen, & Scarpello, 2004). The exploratory bi-factor model was estimated with Bi-Geomin rotation which allowed the sub-factors to correlate with each other. Parallel analysis was done with 1000 randomly generated matrices. The best-fitting exploratory bifactor model was then used to estimate a confirmatory bi-factor model in order to compute the factor scores for subsequent analyses. Model fit indices were also used to evaluate the fit of the exploratory model.

Low factor loadings (i.e., < 0.30) were set to 0 in the confirmatory model. The sub-factors were allowed to correlate with each other but not with the general factor.

All models were estimated using MPLUS 7.4 (Muthén & Muthén, 1998-2012) with maximum likelihood robust estimation.

Correlations. The factor scores derived from the confirmatory bi-factor model were used in the correlation analyses with the child's behavioral outcome measures. Heatmap was plotted to illustrate the patterns of significant correlations between the latent factor scores and outcome measures.

Results

Bi-factor model

Eigenvalues derived from the parallel analysis were used to determine the number of factors. The eigenvalues and fit indices of all the exploratory bi-factor models are summarized in

Table 2. The eigenvalues of seven factors were higher than the randomly generated eigenvalues. Furthermore, the difference in Bayesian Information Criteria (BIC) coefficients of the 7-factor versus 8-factor models was merely 72.75 (0.06%), suggesting little improvement in fit by increasing the number of factors to eight. The comparative fit index (CFI), root mean square error of approximate information (RMSEA) and standardized root mean square residual (SRMR) of the 7-factor model also showed an acceptable fit of data to the model. The general factor explained 62.6% of the common variance extracted and 37.4% were explained by the sub-factors, which corroborated with the results that maternal mental health during pregnancy was multidimensional (Reise, 2012).

The factor loadings of items on the 7-factor model are summarized in Table 3. Factor loadings > 0.30 suggested the item loaded significantly on factor (Hair, Black, Babin, Anderson, & Tatham, 2009). The items that loaded on factors 3 and 7 are of interest to the current study and will be discussed in greater details here. Items that loaded on factor 3 pertained to the STAI items about feeling positive (e.g., feeling pleasant, self-confident, content, satisfied). This factor was thus labelled as Positive Mood. Items that loaded on factor 7 were fewer and pertained to how participants felt or perceived themselves in general (e.g., feeling happy or perceived self as a person who makes decision easily). Factor 7 was labelled as Positive Self. Two items cross-loaded on these two factors. Factors 3 and 7 also had the highest correlation ($r = .24$) as compared to other pairs of factors ($r_s \leq |.15|$).

The 7-factor bi-factor model was estimated using confirmatory bi-factor modelling to obtain the factor scores. The fit indices showed acceptable fit of CFA model to data (RMSEA = .042; CFI = .824; SRMR = .057). Table 4 summarized the items for each factor and the corresponding factor loadings from the confirmatory model.

Reliability indices for the general and sub-factors were also computed (see Table 5) with the excel-based Bifactor Indices Calculator (Dueber, 2016), as previously suggested (Rodriguez,

Reise, & Haviland, 2016). Factor determinacy is the correlation between factor scores and the factors and estimates the reliability the estimated factor score. High factor determinacy coefficient (i.e., ≥ 0.80) suggests high reliability of factor score (Gorsuch, 1983). Both the factor scores of positive mood (FD = .938) and positive self (FD = .89) passed the threshold and were thus used for subsequent analyses. However, according to the Omega Hierarchical coefficient, which reflect the unique variance associated with sub-factor score once partitioning out general factor's variance, Positive mood ($\omega_H = .512$) was more reliable than Positive self ($\omega_H = .427$). The seemingly low omega hierarchical coefficients were not surprising as all the items in these two sub-factors loaded on the general factor as well (Rodriguez et al., 2016), which also accounted for the higher Omega coefficient (ω) that does not control for variance accounted for by general factor. Construct reliability (i.e., *H* index) reflects how well the items represent the latent factor that they load on. With a criterion of 0.70 (Hancock & Mueller, 2001), Positive Mood ($H = .867$) was well-represented by the corresponding sets of items, with Positive Self ($H = .653$) slightly below the threshold. As such, results pertaining to the Positive Self factor score should be interpreted with some caution Nevertheless, these findings suggest that positive mental health construct can be reliably extracted from screening tools for depression and anxiety.

Correlations

The factor scores estimated from the confirmatory model were used in subsequent correlation analyses with the child behavioral outcomes. As this was an exploratory study, Bonferroni correction was not implemented to avoid inflation of Type-II errors. While spurious results may occur without correction, what is of interest is not any particular significant finding, but the pattern of responses which is less likely to be due to chance (Moran, 2003).

The Pearson correlation coefficients and the corresponding p-values are summarized in Table 6. A heatmap (Figure 1) was plotted to better illustrate the pattern of significant

correlations of Positive Mood and Positive Self on child behavioral outcomes. Positive mood and/or self were positively associated with the cognitive, language (i.e., receptive and expressive languages), social-emotional and motor components of the Bayley scales. There were negative association with the total score and social component of the QChat; there was no significant association of positive mood or self on the behavioral component. The positive factors were positively associated with most of the competence subscales of the ITSEA. There was also a positive association of with peer aggression. In general, positive maternal mood during pregnancy were associated with the behaviors in children that are associated with sociability, communication and parentally-rated competence.

Discussion

We used a bi-factor modeling approach to demonstrate the feasibility of using common screening instruments for mental disorders to examine positive maternal mental health. An exploratory analysis revealed associations between our measures of antenatal positive maternal mental health and specific domains of child development. These associations were strongest in measures of social behavior and communication, that were apparent on both maternal report measures as well as those that employ an independent observer (i.e., Bayley Scales). These findings suggest that data from past or existing birth-cohort studies can be examined for potential effects of positive maternal mental health, even in the absence of scales directly intended to examine this construct.

Most of the items on the three mental health questionnaires loaded strongly on the general factor, suggesting an underlying general psychopathology factor that affected the responses of all the items regardless of questionnaires. This could also reflect the comorbidity often found between depression and anxiety measures. Existing research has interpreted this general factor as either general propensity to develop psychopathology symptoms (Caspi et al.,

2014) or general level of distress (Simms et al., 2008). Items about punishment feelings, self-criticalness and loss of interest in sex did not load highly on the general factor.

The sub-factors did not contain items from multiple measures. This finding suggests mental health measures are checklists of different psychopathology symptoms with little overlap. While this may not be surprising as the STAI and BDI are measures of anxiety and depression respectively, there was also no overlap between the two depression measures (BDI and EPDS). Inclusion of multiple mental health measures do have value-add as they may provide a more comprehensive capture of individuals' mental health. This thought is also in line with the understanding that mental health, even within a single mental disorder such as depression, is highly heterogeneous across the population (e.g., Chekroud et al., 2017; Fried, Nesse, Zivin, Guille, & Sen, 2014; Santos et al., 2016).

The positively worded items in the battery loaded negatively on the general factor, suggesting that positive mental health could be a protective factor against psychopathology symptoms. Moreover, these items loaded on two sub-factors that were interpreted as positive mood and positive self, the latter containing less items and is more similar to self-esteem. These two factors are correlated with expected cross-loading of items. The parsing of positively-worded items as distinct factors suggest that it is possible to study positive mental health using commonly used psychiatric screening measures. This finding implies that existing birth cohorts or epidemiological studies with standard screens for maternal psychopathology could be exploited for more comprehensive study of mental health in general population with the existing data used to reveal variations in positive mental health across community samples.

The presence of two separate, but correlated, factors also suggest that positive mental health can be examined from different perspectives. This finding is consistent with existing theoretical conceptualization of psychological well-being that goes beyond merely positive feelings. Psychological well-being has been operationalized into six dimensions: purpose in life,

personal growth, environmental mastery, autonomy, self-acceptance and relations with others (Ryff, 1989). While positive mood is not one of the six traditionally defined dimensions of well-being, it may be a consequence of fulfillment in one or more of the six aspects. Factor analyses of other common psychiatric screening tools have also found a factor of Positive Affect that comprises of the positively worded items (Hernández-Martínez et al., 2011; Hu et al., 2007; Iwata et al., 1998; Shafer, 2006). Deeper research into positive mental health will require a distinction of the different aspects of psychological well-being. However, this does not negate the value of using positive mood, particularly in epidemiological studies that have practical limitations on the measures that can be included.

Effect of positive mental health

Positive antenatal mental health revealed specific associations with child outcomes. Specifically, positive antenatal mental health was significantly associated with cognitive, language/communication, social and competence development. The receptive and expressive language and cognitive abilities were assessed through objective laboratory tasks, which minimize the possibility that parents who were more positive might have rated their child's cognitive and language more positively. Additionally, the parent-rated language component of the QChat had no associations with antenatal maternal mental health. Taken together with the associations with the competence measures, positive antenatal mental health may affect the positive spectrum of a child development instead of socio-emotional vulnerabilities more commonly associated with measures of maternal depression and anxiety. The specificity of the effect of positive mental health is underscored by the finding most of these same measures were not associated with either the sub-factors that reflected a poorer quality of maternal mental health or, in certain instances, even the general factor, despite the liberal p-value threshold used

in this exploratory analysis. Positive maternal mental health may thus have very specific influences on child development.

The pattern of correlations with language, sociability and competences aligns to what is known about children's positive affect, language ability and social traits. Infant positive affect or joyful expressions predict receptive and expressive language abilities in toddlerhood (Dixon Jr & Smith, 2000; Laake & Bridgett, 2014; Moreno & Robinson, 2005). Moreover, a behavioral genetics analysis showed some heritability for sociability and positive affect (Eid, Riemann, Angleitner, & Borkenau, 2003), thus supporting the link between positive affect and sociability.

The effects of positive antenatal mental health on child's language and social abilities may have other long-term indirect benefits. Children who are more sociable may be more accepted by peers, which protects against psychopathological and antisocial behaviors (Parker & Asher, 1987; Szekely et al., 2016). Being more accepted by peers may also contribute to less peer victimization or bullying, which has been found to have serious psychological effects (e.g., Gini & Espelage, 2014; Kawabata, Tseng, & Crick, 2014; Schwartz, Lansford, Dodge, Pettit, & Bates, 2015). These detrimental effects can even last into adulthood as the victimized child enters adolescent and adulthood (McDougall & Vaillancourt, 2015). Taken together with our results, promoting positive antenatal mental health may serve as preventive measures against mental health issues in the next generations.

Promoting positive mental health during pregnancy can also protect against high antenatal stress (see Gaignic-Philippe et al., 2014, for review on antenatal stress and detrimental effects on fetal development). Positive mental health has been associated with better self-care (Giltay, Geleijnse, Zitman, Buijsse, & Kromhout, 2007; Steptoe, Wright, Kunz-Ebrecht, & Iliffe, 2006), higher adherence to medical advice (Cooper, Lloyd, Weinman, & Jackson, 1999), healthier regulation of immune and neuroendocrine systems during stress (Antoni, Carver, & Lechner, 2009; Antoni et al., 2006; Creswell et al., 2005; Sherman, Bunyan, Creswell, &

Jaremka, 2009; Taylor, Lerner, Sherman, Sage, & McDowell, 2003), and lower likelihood of developing clinical depression after experiencing a crisis (Fredrickson, Tugade, Waugh, & Larkin, 2003). Childbirth is a life-changing experience which can be highly stressful or anxiety-provoking, particularly for first-time mothers. Promoting positive antenatal mental health may thus be a more proactive approach to prevent stress-related pregnancy issues before they become severe enough to warrant attention.

Limitations & future research

This exploratory study reveals specific associations between antenatal positive mental health and child development. As such, future research on positive maternal mental health should use measures or laboratory tasks that assess normal development and competences rather than deficit or atypical development. However, current findings are not conclusive and should be interpreted with caution. The mechanisms of how maternal positive mental health affect fetal and child development is unknown. While this study suggests the plausible effects of antenatal positive mental health on fetal and child development, this is an exploratory study with liberal thresholds for significant findings. As such, results require replication.

Another limitation is the lack of postnatal maternal mental health data in this study. As such, we are not able to parcel out the effect of post-natal positive mental health on the child's outcomes. It is possible that antenatal positive mental health persisted into post-natal mental health which in turn affected child outcomes. Nevertheless, this does not negate the importance of antenatal positive mental health. If the effect of antenatal positivity is mediated by post-natal positivity, it then may suggest the importance of intervention or mental health promotion to begin prenatally. Finally, it is important to note the correlational nature of these analyses. This approach cannot discount the possibility of a maternally-inherited effect. Indeed, a recent genome-wide association study (Okbay et al., 2016) described the genetics of emotional well-

being, although genetic variation accounted for only a small percentage of the variation in well-being.

Conclusion

In conclusion, this study demonstrates the feasibility of using common psychiatric disorders screening tools to examine the effect of positive mental health. With this, it is possible for data from past or existing birth-cohort studies to be re-examined from the perspective of positive mental health. Moreover, the effects of positive mental health are likely to be specific and different from the lack of mental disorders. As such, a deeper understanding of positive mental health will allow for more comprehensive understanding of fetal and child development. This also highlights the need to promote mental health among the general population in addition to preventing mental disorders.

Table 1
Contents of all items used in the bi-factor exploratory analysis.

| Questionnaire | Item | Content |
|---|--------|---|
| Edinburgh Postnatal Depression Scale | EPDS1 | In the past 7 days, I have been able to laugh and see the funny side of things. |
| | EPDS2 | In the past 7 days, I have looked forward with enjoyment to things. |
| | EPDS3 | In the past 7 days, I have blamed myself unnecessarily when things went wrong. |
| | EPDS4 | In the past 7 days, I have been anxious or worried for no good reason. |
| | EPDS5 | In the past 7 days, I have felt scared or panicky for no very good reason. |
| | EPDS6 | In the past 7 days, things have been getting on top of me. |
| | EPDS7 | In the past 7 days, I have been so unhappy that I have had difficulty sleeping. |
| | EPDS8 | In the past 7 days, I have felt sad or miserable. |
| | EPDS9 | In the past 7 days, I have been so unhappy that I have been crying. |
| | EPDS10 | In the past 7 days, the thought of harming myself has occurred to me. |
| Beck Depression Inventory-II | BDI1 | Sadness |
| | BDI2 | Pessimism |
| | BDI3 | Past Failure |
| | BDI4 | Loss of Pleasure |
| | BDI5 | Guilty Feelings |
| | BDI6 | Punishment Feelings |
| | BDI7 | Self-Dislike |
| | BDI8 | Self-Criticalness |
| | BDI9 | Suicidal Thoughts or Wishes |
| | BDI10 | Crying |
| | BDI11 | Agitation |
| | BDI12 | Loss of Interest |
| | BDI13 | Indecisiveness |
| | BDI14 | Worthlessness |
| | BDI15 | Loss of Energy |
| | BDI16 | Changes in Sleeping Pattern |
| | BDI17 | Irritability |
| | BDI18 | Changes in Appetite |
| | BDI19 | Concentration Difficulty |
| | BDI20 | Tiredness or Fatigue |
| | BDI21 | Loss of Interest in Sex |
| State-Trait Anxiety Inventory | STAI1 | I feel calm (at this moment) |
| | STAI2 | I feel secure (at this moment) |
| | STAI3 | I am tense (at this moment) |
| | STAI4 | I feel strained (at this moment) |
| | STAI5 | I feel at ease (at this moment) |
| | STAI6 | I feel upset (at this moment) |
| | STAI7 | I am presently worrying over possible misfortunes (at this moment) |

| Questionnaire | Item | Content |
|---------------|--------|---|
| | STAI8 | I feel satisfied (at this moment) |
| | STAI9 | I feel frightened (at this moment) |
| | STAI10 | I feel comfortable (at this moment) |
| | STAI11 | I feel self-confident (at this moment) |
| | STAI12 | I feel nervous (at this moment) |
| | STAI13 | I am jittery (at this moment) |
| | STAI14 | I feel indecisive (at this moment) |
| | STAI15 | I am relaxed (at this moment) |
| | STAI16 | I feel content (at this moment) |
| | STAI17 | I am worried (at this moment) |
| | STAI18 | I feel confused (at this moment) |
| | STAI19 | I feel steady (at this moment) |
| | STAI20 | I feel pleasant (at this moment) |
| | STAI21 | I feel pleasant (generally) |
| | STAI22 | I feel nervous and restless (generally) |
| | STAI23 | I feel satisfied with myself (generally) |
| | STAI24 | I wish I could be as happy as others seem to be (generally) |
| | STAI25 | I feel like a failure (generally) |
| | STAI26 | I feel rested (generally) |
| | STAI27 | I am calm, cool, and collected (generally) |
| | STAI28 | I feel that difficulties are piling up so that I cannot overcome them (generally) |
| | STAI29 | I worry too much over something that really doesn't matter (generally) |
| | STAI30 | I am happy (generally) |
| | STAI31 | I have disturbing thoughts (generally) |
| | STAI32 | I lack self-confidence (generally) |
| | STAI33 | I feel secure (generally) |
| | STAI34 | I make decisions easily (generally) |
| | STAI35 | I feel inadequate (generally) |
| | STAI36 | I am content (generally) |
| | STAI37 | Some unimportant thought runs through my mind and bothers me (generally) |
| | STAI38 | I take disappointments so keenly that I can't put them out of my mind (generally) |
| | STAI39 | I am a steady person (generally) |
| | STAI40 | I get in a state of tension or turmoil as I think over my recent concerns and interests (generally) |

Table 2
Bi-factor model fit statistics

| Model | Eigenvalue | Eigenvalue from parallel analysis | Model df | χ^2 | AIC | BIC | Adjusted BIC | RMSEA | RMSEA p-value | CFI | SRMR |
|-----------------------|------------|--|-------------|----------|-----------|-----------|-----------------|-------|------------------|------|------|
| Exploratory models | | | | | | | | | | | |
| 1-factor ^a | 19.12 | 1.55 | 2345 | 14781.11 | 135065.95 | 136110.00 | 135443.00 | .071 | <.001 | .609 | .077 |
| 2-factor | 4.64 | 1.51 | 2276 | 10025.40 | 130448.24 | 131835.34 | 130949.18 | .057 | <.001 | .757 | .052 |
| 3-factor | 3.00 | 1.48 | 2208 | 8165.56 | 128724.40 | 130449.57 | 129347.43 | .050 | .329 | .813 | .041 |
| 4-factor | 2.11 | 1.46 | 2141 | 7279.17 | 127972.01 | 130030.28 | 128715.34 | .047 | 1.00 | .839 | .036 |
| 5-factor | 1.81 | 1.43 | 2075 | 6434.20 | 127259.04 | 129645.44 | 128120.87 | .044 | 1.00 | .863 | .032 |
| 6-factor | 1.61 | 1.40 | 2010 | 5703.83 | 126658.67 | 129368.22 | 127637.21 | .042 | 1.00 | .884 | .029 |
| 7-factor | 1.43 | 1.39 | 1946 | 5134.65 | 126217.49 | 129245.23 | 127310.94 | .039 | 1.00 | .900 | .027 |
| 8-factor | 1.33 | 1.36 | 1883 | 4622.68 | 125831.52 | 129172.48 | 127038.09 | .037 | 1.00 | .914 | .025 |
| Confirmatory model | | | | | | | | | | | |
| 7-factor | - | - | 2359 | 6809.58 | 131018.16 | 132350.57 | 131499.354 | .042 | 1.00 | .824 | .057 |

^aThis is a unidimensional model, not a bi-factor model that requires at least 2 factors (1 general & 1 specific).

Note. AIC = Akaike Information Criteria; BIC = Bayesian Information Criteria; RMSEA = root mean square error of approximation; CFI = comparative fit index; SRMR = standardized root mean square residual.

Table 3
Factor loadings & correlations from 7-factor exploratory bifactor model

| Item | G-factor | Factor 2 | Factor 3 | Factor 4 | Factor 5 | Factor 6 | Factor 7 |
|--------|--------------|-------------|-------------|-------------|--------------|-------------|----------|
| BDI1 | .548 | .177 | .043 | .004 | -.215 | .064 | .005 |
| BDI2 | .418 | .258 | -.004 | -.051 | -.104 | -.010 | .003 |
| BDI3 | .474 | .334 | .068 | -.016 | .027 | -.068 | -.025 |
| BDI4 | .494 | .159 | -.04 | .183 | -.092 | .059 | -.039 |
| BDI5 | .499 | .290 | .062 | .023 | -.045 | -.050 | .040 |
| BDI6 | .363 | .293 | .07 | -.102 | -.012 | -.008 | .031 |
| BDI7 | .431 | .389 | -.027 | .092 | .023 | -.084 | .069 |
| BDI8 | .481 | .328 | .037 | -.033 | -.03 | -.080 | .009 |
| BDI9 | .336 | .425 | -.026 | .046 | -.023 | -.024 | .017 |
| BDI10 | .52 | .115 | .08 | .047 | -.223 | -.003 | -.031 |
| BDI11 | .484 | .144 | .043 | .240 | -.043 | .036 | -.042 |
| BDI12 | .469 | .177 | -.003 | .275 | -.015 | .039 | -.055 |
| BDI13 | .502 | .230 | .052 | .191 | .066 | -.017 | -.097 |
| BDI14 | .51 | .446 | .002 | .057 | .031 | -.045 | .039 |
| BDI15 | .359 | -.123 | .031 | .591 | -.024 | .013 | -.028 |
| BDI16 | .274 | -.005 | -.008 | .366 | -.02 | -.035 | .083 |
| BDI17 | .473 | .075 | .02 | .352 | .002 | -.050 | .011 |
| BDI18 | .139 | .040 | 0 | .334 | -.065 | -.020 | -.001 |
| BDI19 | .478 | .057 | -.007 | .385 | .07 | -.047 | -.023 |
| BDI20 | .329 | -.028 | -.01 | .551 | .026 | -.016 | .083 |
| BDI21 | .154 | .043 | -.054 | .262 | -.012 | 0 | .049 |
| EPDS1 | -.343 | -.232 | .098 | -.138 | .226 | -.085 | .028 |
| EPDS2 | -.384 | -.169 | .136 | -.146 | .172 | -.029 | .079 |
| EPDS3 | .500 | -.085 | -.008 | -.197 | -.024 | -.171 | .004 |
| EPDS4 | .565 | -.300 | .102 | .023 | -.057 | -.164 | -.019 |
| EPDS5 | .565 | -.191 | .048 | -.041 | -.093 | -.082 | .058 |
| EPDS6 | .520 | -.176 | .005 | .075 | -.107 | -.147 | .069 |
| EPDS7 | .567 | -.035 | -.042 | -.025 | -.298 | -.015 | .135 |
| EPDS8 | .678 | -.06 | .034 | -.047 | -.372 | -.040 | .015 |
| EPDS9 | .587 | -.024 | .042 | -.098 | -.360 | -.071 | .034 |
| EPDS10 | .472 | .181 | .019 | -.146 | -.203 | -.058 | .059 |
| STAI1 | -.454 | .009 | .518 | -.011 | .06 | -.081 | -.044 |
| STAI2 | -.460 | -.057 | .588 | .042 | -.032 | -.026 | -.030 |
| STAI3 | .443 | -.054 | -.114 | .034 | -.054 | .397 | .150 |
| STAI4 | .384 | -.107 | -.058 | .047 | -.158 | .432 | .100 |
| STAI5 | -.425 | .027 | .594 | .035 | -.055 | .002 | -.210 |
| STAI6 | .555 | .058 | -.059 | -.055 | -.061 | .364 | .017 |
| STAI7 | .494 | -.023 | .040 | .014 | .082 | .322 | -.015 |
| STAI8 | -.430 | .002 | .560 | .023 | -.048 | .001 | -.007 |
| STAI9 | .477 | .048 | .052 | -.075 | .021 | .378 | -.043 |
| STAI10 | -.427 | .022 | .560 | -.034 | .032 | -.030 | .059 |
| STAI11 | -.442 | .019 | .582 | -.061 | -.124 | .048 | .054 |
| STAI12 | .517 | -.02 | .075 | .034 | .046 | .423 | .018 |
| STAI13 | .485 | -.024 | .004 | -.049 | .205 | .360 | .035 |
| STAI14 | .448 | -.01 | .004 | -.001 | .251 | .235 | .009 |
| STAI15 | -.498 | .006 | .615 | -.072 | -.053 | -.020 | -.029 |
| STAI16 | -.435 | -.019 | .600 | .036 | .105 | .052 | -.088 |

| | | | | | | | |
|---------------------|--------------|-------|-------------|-------|-------|-------------|-------------|
| STAI17 | .567 | -.036 | .07 | -.015 | .037 | .395 | -.144 |
| STAI18 | .554 | .067 | .057 | -.027 | .035 | .412 | -.078 |
| STAI19 | -.476 | -.06 | .604 | -.008 | .023 | .031 | .016 |
| STAI20 | -.479 | -.055 | .643 | -.023 | .003 | .001 | .011 |
| STAI21 | -.535 | -.021 | .410 | .018 | .122 | -.053 | .333 |
| STAI22 | .605 | -.089 | .004 | .056 | .162 | .156 | .069 |
| STAI23 | -.533 | -.025 | .390 | -.019 | -.061 | .018 | .354 |
| STAI24 | .290 | .082 | .020 | -.085 | .012 | .055 | .181 |
| STAI25 | .536 | .206 | -.026 | -.188 | .25 | .016 | -.031 |
| STAI26 | -.380 | .105 | .301 | -.071 | -.007 | .008 | .142 |
| STAI27 | -.566 | .046 | .364 | -.005 | -.073 | .067 | .273 |
| STAI28 | .594 | .028 | -.022 | -.019 | .256 | -.001 | .099 |
| STAI29 | .603 | -.195 | .110 | -.021 | .063 | -.008 | -.067 |
| STAI30 | -.531 | .046 | .280 | .002 | .197 | -.022 | .364 |
| STAI31 | .648 | -.024 | .047 | -.052 | .194 | .104 | .029 |
| STAI32 | .536 | .082 | .051 | -.035 | .196 | .003 | -.085 |
| STAI33 | -.505 | .011 | .425 | .110 | .003 | -.020 | .245 |
| STAI34 | -.536 | .042 | .215 | -.023 | -.156 | .141 | .311 |
| STAI35 | .478 | -.018 | -.012 | -.111 | .297 | .037 | .048 |
| STAI36 | -.513 | .011 | .421 | .099 | .125 | .008 | .123 |
| STAI37 | .584 | -.148 | .082 | .079 | .052 | .056 | .056 |
| STAI38 | .607 | -.068 | .081 | .092 | .04 | .093 | .019 |
| STAI39 | -.549 | -.026 | .306 | .005 | .028 | .041 | .360 |
| STAI40 | .580 | -.076 | -.037 | -.058 | .238 | .073 | .173 |
| Factor correlations | | | | | | | |
| Factor 2 | .00 | 1 | | | | | |
| Factor 3 | .00 | -.02 | 1 | | | | |
| Factor 4 | .00 | .11* | -.03 | 1 | | | |
| Factor 5 | .00 | -.09* | .04 | -.07* | 1 | | |
| Factor 6 | .00 | -.12* | .03 | -.03 | .15* | 1 | |
| Factor 7 | .00 | -.06 | .22 | -.07* | -.02 | .10 | 1 |

Note. Bold indicates factor loading > |.30|. *significant correlation at $p < .05$.

Table 4
Specific latent factors & contents of items

| EFA factor loadings | | Item | Contents |
|--------------------------|------------|--------|---|
| General factor | Sub-factor | | |
| Factor 2 (Self-loath) | | | |
| 0.51 | 0.47 | BDI14 | Worthlessness |
| 0.34 | 0.43 | BDI9 | Suicidal Thoughts or Wishes |
| 0.43 | 0.39 | BDI7 | Self-Dislike |
| 0.47 | 0.33 | BDI3 | Past Failure |
| 0.48 | 0.33 | BDI8 | Self-Criticalness |
| Factor 3 (Positive mood) | | | |
| -0.48 | 0.64 | STAI20 | I feel pleasant (at this moment) |
| -0.48 | 0.60 | STAI19 | I feel steady (at this moment) |
| -0.44 | 0.60 | STAI16 | I feel content (at this moment) |
| -0.43 | 0.59 | STAI5 | I feel at ease (at this moment) |
| -0.46 | 0.59 | STAI2 | I feel secure (at this moment) |
| -0.44 | 0.58 | STAI11 | I feel self-confident (at this moment) |
| -0.43 | 0.59 | STAI15 | I am relaxed (at this moment) |
| -0.43 | 0.56 | STAI10 | I feel satisfied (at this moment) |
| -0.43 | 0.56 | STAI8 | I feel comfortable (at this moment) |
| -0.45 | 0.52 | STAI1 | I feel calm (at this moment) |
| -0.51 | 0.43 | STAI33 | I feel secure (generally) |
| -0.51 | 0.42 | STAI36 | I am content (generally) |
| -0.54 | 0.41 | STAI21 | I feel pleasant (generally) ^a |
| -0.53 | 0.39 | STAI23 | I feel satisfied with myself (generally) ^a |
| -0.57 | 0.36 | STAI27 | I am calm, cool, and collected (generally) |
| -0.55 | 0.21 | STAI39 | I am a steady person (generally) ^a |
| -0.38 | 0.30 | STAI26 | I feel rested (generally) |
| Factor 4 (Psychosomatic) | | | |
| 0.34 | 0.59 | BDI15 | Loss of Energy |
| 0.33 | 0.55 | BDI20 | Tiredness or Fatigue |
| 0.48 | 0.39 | BDI19 | Concentration Difficulty |
| 0.27 | 0.37 | BDI16 | Changes in Sleeping Pattern |
| 0.47 | 0.35 | BDI17 | Irritability |
| 0.14 | 0.33 | BDI18 | Changes in Appetite |
| Factor 5 (Melancholy) | | | |
| 0.68 | 0.37 | EPDS8 | In the past 7 days, I have felt sad or miserable. |
| 0.59 | 0.36 | EPDS9 | In the past 7 days, I have been so unhappy that I have been crying. |
| Factor 6 (Anxiety) | | | |
| 0.38 | 0.43 | STAI4 | I feel strained (at this moment) |
| 0.52 | 0.42 | STAI12 | I feel nervous (at this moment) |
| 0.55 | 0.41 | STAI18 | I feel confused (at this moment) |
| 0.44 | 0.40 | STAI3 | I am tense (at this moment) |
| 0.57 | 0.40 | STAI17 | I am worried (at this moment) |
| 0.48 | 0.38 | STAI9 | I feel frightened (at this moment) |
| 0.56 | 0.36 | STAI6 | I feel upset (at this moment) |
| 0.49 | 0.36 | STAI13 | I am jittery (at this moment) |

| EFA factor loadings | | | |
|--------------------------|------------|--------|---|
| General factor | Sub-factor | Item | Contents |
| 0.49 | 0.32 | STAI7 | I am presently worrying over possible misfortunes. |
| Factor 7 (Positive self) | | | |
| -0.53 | 0.36 | STAI30 | I am happy (generally) |
| -0.55 | 0.36 | STAI39 | I am a steady person (generally) ^a |
| -0.53 | 0.35 | STAI23 | I feel satisfied with myself (generally) ^a |
| -0.54 | 0.33 | STAI21 | I feel pleasant (generally) ^a |
| -0.54 | 0.31 | STAI34 | I make decisions easily (generally) |

^a item in cross-loaded on factors 3 & 7.

Table 5
Reliability indices from confirmatory bi-factor model

| Factors | Factor determinacy | ω | ω_H | <i>H</i> index |
|---------------|--------------------|----------|------------|----------------|
| General | .970 | .861 | .581 | .957 |
| Self-loath | .778 | .475 | .344 | .544 |
| Positive mood | .938 | .575 | .512 | .867 |
| Psychosomatic | .820 | .715 | .553 | .652 |
| Melancholy | .761 | .405 | .301 | .425 |
| Anxiety | .824 | .378 | .325 | .626 |
| Positive self | .890 | .504 | .427 | .653 |

Note. ω = omega coefficient. ω_H = omega hierarchical coefficient. *H* = construct reliability.

Table 6.
Pearson correlations & p-values (in italics) of latent factor scores and child behavioral outcomes

| Measure | Subscales | Latent factor scores | | | | | | |
|--|--------------------------|-----------------------|----------------------|-----------------------------|-----------------------------|----------------------|----------------------|-----------------------------|
| | | General | Loath | Positive Mood | Psychosomatic | Melancholy | Anxiety | Positive Self |
| Bayley Scale of Infant & Toddler Development | Cognitive | -.034 <i>.467</i> | -.013 <i>.781</i> | 0.133 <i>.005</i> | 0.137 <i>.003</i> | -.049 <i>.299</i> | -.048 <i>.306</i> | 0.107 <i>.023</i> |
| | Receptive communication | -.091 <i>.053</i> | .019 <i>.682</i> | 0.115 <i>.015</i> | .052 <i>.267</i> | -.025 <i>.603</i> | -.082 <i>.082</i> | .088 <i>.062</i> |
| | Expressive communication | -.084 <i>.077</i> | .012 <i>.805</i> | 0.133 <i>.005</i> | .084 <i>.075</i> | -.052 <i>.270</i> | -.073 <i>.122</i> | .091 <i>.054</i> |
| | Fine motor skills | -.070 <i>.139</i> | -.063 <i>.183</i> | .047 <i>.318</i> | .066 <i>.163</i> | -.058 <i>.219</i> | -.012 <i>.799</i> | .034 <i>.472</i> |
| | Gross motor skills | -.004 <i>.928</i> | -.061 <i>.199</i> | .064 <i>.178</i> | .083 <i>.080</i> | .040 <i>.403</i> | -.004 <i>.930</i> | .049 <i>.301</i> |
| | Social-emotional | -0.186 <i>.000</i> | -.053 <i>.280</i> | 0.136 <i>.005</i> | .003 <i>.951</i> | -.085 <i>.080</i> | -.076 <i>.117</i> | 0.099 <i>.041</i> |
| | Communication behavior | -.038 <i>.437</i> | .055 <i>.260</i> | .057 <i>.239</i> | -.048 <i>.324</i> | -.082 <i>.090</i> | .029 <i>.547</i> | .057 <i>.239</i> |
| | Community use | -.037 <i>.450</i> | .017 <i>.727</i> | .001 <i>.985</i> | -.075 <i>.121</i> | -.045 <i>.358</i> | .000 <i>.996</i> | -.023 <i>.630</i> |
| | Functional pre-academic | -.015 <i>.760</i> | -.022 <i>.653</i> | .092 <i>.057</i> | -.006 <i>.909</i> | -.031 <i>.524</i> | -.015 <i>.763</i> | .086 <i>.076</i> |
| | Home living | -.070 <i>.150</i> | .031 <i>.518</i> | .010 <i>.840</i> | .007 <i>.887</i> | -.053 <i>.272</i> | .011 <i>.817</i> | .006 <i>.905</i> |

| Measure | Subscales | Latent factor scores | | | | | | |
|--|---|-----------------------|---------------|-----------------------|---------------|---------------|---------------|-----------------------|
| | | General | Loath | Positive Mood | Psychosomatic | Melancholy | Anxiety | Positive Self |
| | Health & safety knowledge | -0.142 .003 | .073 .133 | .090 .066 | -.031 .521 | -.080 .099 | -.069 .153 | .052 .284 |
| | Leisure activities | -0.103 .033 | .055 .259 | .090 .063 | -.020 .681 | -.080 .098 | -.088 .069 | .058 .230 |
| | Self care | -.062 .201 | .068 .159 | .025 .607 | -.019 .695 | -.042 .392 | -.008 .871 | .017 .721 |
| | Self-direction skills | -0.1 .038 | .067 .168 | .010 .842 | -.045 .360 | -.058 .233 | -.017 .721 | .003 .959 |
| | Social skills | -0.122 .012 | .006 .905 | .045 .357 | -.068 .163 | -.093 .054 | -.033 .499 | .031 .523 |
| | Motor skills | -0.135 .005 | -.076 .120 | 0.111 .022 | -.028 .559 | -.095 .051 | .015 .763 | 0.11 .023 |
| Qchat 18months | Total score | 0.171 .001 | -.075 .163 | -0.142 .008 | -.022 .686 | .014 .794 | .060 .265 | -0.132 .014 |
| | Social factor | .052 .332 | .040 .459 | -0.16 .003 | .044 .414 | -.012 .828 | .043 .427 | -0.149 .006 |
| | Behavioral factor | 0.16 .003 | -.072 .181 | -.040 .457 | -.045 .404 | -.003 .957 | .037 .492 | -.050 .353 |
| | Language factor | .060 .268 | -.051 .342 | -.033 .543 | -.028 .608 | .026 .625 | .056 .298 | -.038 .477 |
| Infant Toddler Social-Emotional Assessment | Activity/Impulsivity (externalizing) | 0.143 .003 | .017 .713 | .010 .838 | .063 .182 | -.011 .820 | .025 .598 | .021 .654 |
| | Aggression/Defiance (externalizing) | 0.167 .000 | .006 .894 | .001 .986 | .050 .298 | -.070 .145 | .079 .097 | .033 .488 |

| Measure | Subscales | Latent factor scores | | | | | | |
|--|-----------------------|----------------------|----------------------|---------------|-----------------------|----------------------|----------------------|---------------|
| | | General | Loath | Positive Mood | Psychosomatic | Melancholy | Anxiety | Positive Self |
| Peer Aggression (externalizing) | 0.115 .032 | .074 .166 | 0.132 .013 | .068 .203 | -0.136 .010 | .064 .228 | 0.142 .008 | |
| Depression/Withdrawal (internalizing) | 0.225 .000 | .051 .288 | -.021 .652 | -.028 .556 | -.027 .566 | .090 .058 | .023 .636 | |
| Fear | .080 .098 | -.001 .990 | .006 .900 | -.022 .653 | -.018 .705 | .025 .602 | .034 .482 | |
| General anxiety (internalizing) | 0.179 .000 | .017 .726 | -.031 .516 | -.076 .108 | .012 .809 | .090 .058 | .059 .215 | |
| Separation distress (Internalizing) | .087 .068 | .024 .609 | -.043 .368 | .051 .283 | .028 .563 | -.053 .263 | -.020 .675 | |
| Inhibition to novelty (internalizing) | .048 .319 | -.003 .952 | .032 .500 | .019 .687 | -.032 .507 | -.012 .808 | .053 .272 | |
| Negative emotionality (dysregulation) | 0.257 .000 | .083 .081 | -.001 .975 | .047 .324 | .003 .950 | .009 .851 | .026 .584 | |
| Sleep dysregulation | .079 .097 | -.081 .091 | .032 .502 | .031 .521 | -.009 .854 | .069 .148 | .046 .339 | |
| Eating dysregulation | 0.13 .007 | -.002 .970 | -.059 .218 | .054 .257 | .006 .904 | 0.098 .041 | -.042 .376 | |
| Sensory sensitivity (dysregulation) | 0.112 .019 | .025 .604 | .047 .322 | .052 .276 | .003 .944 | 0.105 .028 | 0.118 .013 | |
| Compliance (competence) | -0.118 .013 | .003 .949 | 0.115 .015 | -.068 .154 | -.059 .220 | .025 .606 | 0.133 .005 | |

| Measure | Subscales | Latent factor scores | | | | | | |
|---------|---------------------------------------|-----------------------|---------------|----------------------|-----------------------|---------------|---------------|----------------------|
| | | General | Loath | Positive Mood | Psychosomatic | Melancholy | Anxiety | Positive Self |
| | Attention (competence) | -0.11 .021 | -.020 .677 | 0.15 .002 | -.027 .569 | -.066 .163 | .091 .057 | 0.178 .000 |
| | Mastery motivation (competence) | -.091 .056 | .064 .178 | 0.119 .012 | .038 .424 | -.021 .665 | .025 .592 | 0.142 .003 |
| | Imitation/Play (competence) | -.055 .248 | .045 .346 | 0.101 .034 | -.009 .858 | -.049 .302 | -.004 .932 | 0.115 .016 |
| | Empathy (competence) | -.003 .943 | .046 .341 | 0.123 .010 | -.073 .126 | -.048 .315 | .093 .052 | 0.144 .002 |
| | Prosocial peer relations (competence) | -.046 .383 | -.009 .862 | -.021 .686 | -0.164 .002 | -.055 .297 | .050 .342 | .017 .754 |
| | Maladaptive item cluster | .087 .068 | -.079 .099 | -.049 .311 | -.001 .987 | -.039 .421 | .093 .051 | .004 .933 |
| | Social relatedness item cluster | -0.146 .002 | .042 .382 | 0.094 .047 | .061 .200 | -.083 .081 | -.053 .267 | 0.108 .023 |
| | Atypical item cluster | 0.128 .007 | -.063 .185 | -.090 .059 | .026 .592 | -.005 .910 | .065 .177 | -.037 .441 |

Note. Bold indicates significant correlation at $\alpha = .05$ level.




Figure 1. Heatmap illustrating significant correlations between maternal mental health factors & child behavioral outcomes.

References

- Allison, C., Baron-Cohen, S., Wheelwright, S., Charman, T., Richler, J., Pasco, G., & Brayne, C. (2008). The Q-CHAT (Quantitative CHECKlist for Autism in Toddlers): a normally distributed quantitative measure of autistic traits at 18–24 months of age: preliminary report. *Journal of Autism and Developmental Disorders*, *38*(8), 1414-1425. doi:10.1007/s10803-007-0509-7
- Antoni, M. H., Carver, C. S., & Lechner, S. C. (2009). Enhancing positive adaptation: Example intervention during treatment for breast cancer. In C. L. Park, S. C. Lechner, M. H. Antoni, & A. L. Stanton (Eds.), *Medical Illness and Positive Life Change* (pp. 197-214). Washington: USA: Psychological Association.
- Antoni, M. H., Lutgendorf, S. K., Cole, S. W., Dhabhar, F. S., Sephton, S. E., McDonald, P. G., . . . Sood, A. K. (2006). The influence of bio-behavioural factors on tumour biology: Pathways and mechanisms. *Nature Reviews Cancer*, *6*(3), 240. doi:10.1038/nrc1820
- Aspinwall, L. G., & Tedeschi, R. G. (2010). The value of positive psychology for health psychology: Progress and pitfalls in examining the relation of positive phenomena to health. *Annals of Behavioral Medicine*, *39*(1), 4-15. doi:10.1007/s12160-009-9153-0
- Babenko, O., Kovalchuk, I., & Metz, G. A. (2015). Stress-induced perinatal and transgenerational epigenetic programming of brain development and mental health. *Neuroscience & Biobehavioral Reviews*, *48*, 70-91. doi:10.1016/j.neubiorev.2014.11.013
- Baibazarova, E., van de Beek, C., Cohen-Kettenis, P. T., Buitelaar, J., Shelton, K. H., & van Goozen, S. H. M. (2013). Influence of prenatal maternal stress, maternal plasma cortisol and cortisol in the amniotic fluid on birth outcomes and child temperament at 3 months. *Psychoneuroendocrinology*, *38*(6), 907-915. doi:10.1016/j.psyneuen.2012.09.015
- Bayley, N. (2006). *Bayley Scales of Infant and Toddler Development—Third Edition*. San Antonio, TX: Harcourt Assessment.

- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Bennett, H. A., Einarson, A., Taddio, A., Koren, G., & Einarson, T. R. (2004). Prevalence of depression during pregnancy: Systematic review. *Obstetrics and Gynecology*, *103*(4), 698-709. doi:10.1097/01.AOG.0000116689.75396.5f
- Beydoun, H., & Saftlas, A. F. (2008). Physical and mental health outcomes of prenatal maternal stress in human and animal studies: a review of recent evidence. *Paediatric and Perinatal Epidemiology*, *22*(5), 438-466. doi:10.1111/j.1365-3016.2008.00951.x
- Bowers, M. E., & Yehuda, R. (2016). Intergenerational transmission of stress in humans. *Neuropsychopharmacology*, *41*(1), 232-244. doi:10.1038/npp.2015.247
- Briggs-Gowan, M. J., & Carter, A. S. (1998). Preliminary acceptability and psychometrics of the infant-toddler social and emotional assessment (ITSEA): A new adult-report questionnaire. *Infant Mental Health Journal*, *19*(4), 422-445. doi:10.1002/(SICI)1097-0355(199824)19:4<422::AID-IMHJ5>3.0.CO;2-U
- Buss, C., Davis, E. P., Shahbaba, B., Pruessner, J. C., Head, K., & Sandman, C. A. (2012). Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumes and affective problems. *Proceedings of the National Academy of Sciences*, *109*(20), E1312-E1319. doi:10.1073/pnas.1201295109
- Caspi, A., Houts, R. M., Belsky, D. W., Goldman-Mellor, S. J., Harrington, H., Israel, S., . . . Poulton, R. (2014). The p factor: One general psychopathology factor in the structure of psychiatric disorders? *Clinical Psychological Science*, *2*(2), 119-137. doi:10.1177/2167702613497473
- Castro, R. T. A., Anderman, C. P., Glover, V., O'Connor, T. G., Ehlert, U., & Kammerer, M. (2016). Associated symptoms of depression: patterns of change during pregnancy. *Archives of Women's Mental Health*, 1-6. doi:10.1007/s00737-016-0685-6

Charil, A., Laplante, D. P., Vaillancourt, C., & King, S. (2010). Prenatal stress and brain development. *Brain Research Reviews*, *65*(1), 56-79.

doi:10.1016/j.brainresrev.2010.06.002

Chekroud, A. M., Gueorguieva, R., Krumholz, H. M., Trivedi, M. H., Krystal, J. H., & McCarthy, G. (2017). Reevaluating the efficacy and predictability of antidepressant treatments: a symptom clustering approach. *Jama psychiatry*, *74*(4), 370-378.

doi:10.1001/jamapsychiatry.2017.0025

Cooper, A., Lloyd, G., Weinman, J., & Jackson, G. (1999). Why patients do not attend cardiac rehabilitation: role of intentions and illness beliefs. *Heart*, *82*(2), 234-236.

doi:10.1136/hrt.82.2.234

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression.

Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal of Psychiatry*, *150*(6), 782-786. doi:10.1192/bjp.150.6.782

Creswell, J. D., Welch, W. T., Taylor, S. E., Sherman, D. K., Gruenewald, T. L., & Mann, T. (2005). Affirmation of personal values buffers neuroendocrine and psychological stress responses. *Psychological Science*, *16*(11), 846-851. doi:10.1111/j.1467-

9280.2005.01624.x

Danner, D. D., Snowdon, D. A., & Friesen, W. V. (2001). Positive emotions in early life and longevity: Findings from the nun study. *Journal of Personality and Social Psychology*, *80*(5), 804-813. doi: 10.1037//0022-3514.80.5.804

de Cates, A., Stranges, S., Blake, A., & Weich, S. (2015). Mental well-being: An important outcome for mental health services? *The British Journal of Psychiatry*, *207*(3), 195-197.

doi:10.1192/bjp.bp.114.158329

Dixon Jr, W. E., & Smith, P. H. (2000). Links between early temperament and language acquisition. *Merrill-Palmer Quarterly*, *46*(3), 417-440.

- Dueber, D. M. (2016). Bifactor Indices Calculator: A Microsoft Excel-based tool to calculate various indices relevant to bifactor CFA models. Retrieved from <http://sites.education.uky.edu/apslab/resources>
- Eid, M., Riemann, R., Angleitner, A., & Borkenau, P. (2003). Sociability and positive emotionality: Genetic and environmental contributions to the covariation between different facets of extraversion. *Journal of Personality, 71*(3), 319-346.
- Field, T. (2011). Prenatal depression effects on early development: A review. *Infant Behavior and Development, 34*(1), 1-14. doi:10.1016/j.infbeh.2010.09.008
- Fredrickson, B. L., Tugade, M. M., Waugh, C. E., & Larkin, G. R. (2003). What good are positive emotions in crisis? A prospective study of resilience and emotions following the terrorist attacks on the United States on September 11th, 2001. *Journal of Personality and Social Psychology, 84*(2), 365. doi:10.1037/0022-3514.84.2.365
- Fried, E. I. (2017). The 52 symptoms of major depression: Lack of content overlap among seven common depression scales. *Journal of Affective Disorders, 208*, 191-197. doi:10.1016/j.jad.2016.10.019
- Fried, E. I., Nesse, R. M., Zivin, K., Guille, C., & Sen, S. (2014). Depression is more than the sum score of its parts: Individual DSM symptoms have different risk factors. *Psychological Medicine, 44*(10), 2067-2076. doi:10.1017/S0033291713002900
- Gavin, N. I., Gaynes, B. N., Lohr, K. N., Meltzer-Brody, S., Gartlehner, G., & Swinson, T. (2005). Perinatal depression: A systematic review of prevalence and incidence. *Obstetrics and Gynecology, 106*(5, Part 1), 1071-1083. doi:10.1097/01.AOG.0000183597.31630.db
- Gentile, S. (2017). Untreated depression during pregnancy: Short-and long-term effects in offspring. A systematic review. *Neuroscience, 342*, 154-166. doi:10.1016/j.neuroscience.2015.09.001

- Giltay, E. J., Geleijnse, J. M., Zitman, F. G., Buijsse, B., & Kromhout, D. (2007). Lifestyle and dietary correlates of dispositional optimism in men: The Zutphen Elderly Study. *Journal of Psychosomatic Research, 63*(5), 483-490. doi:10.1016/j.jpsychores.2007.07.014
- Gini, G., & Espelage, D. L. (2014). Peer victimization, cyberbullying, and suicide risk in children and adolescents. *JAMA, 312*(5), 545-546. doi:10.1001/jama.2014.3212
- Glover, V. (2014). Maternal depression, anxiety and stress during pregnancy and child outcome; what needs to be done. *Best practice & research Clinical obstetrics & gynaecology, 28*(1), 25-35. doi:10.1016/j.bpobgyn.2013.08.017
- Goodman, S. H., Rouse, M. H., Connell, A. M., Broth, M. R., Hall, C. M., & Heyward, D. (2011). Maternal depression and child psychopathology: A meta-analytic review. *Clinical Child and Family Psychology Review, 14*(1), 1-27. doi:10.1007/s10567-010-0080-1
- Gorsuch, R. L. (1983). *Factor analysis* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum Associates.
- Graignic-Philippe, R., Dayan, J., Chokron, S., Jacquet, A. Y., & Tordjman, S. (2014). Effects of prenatal stress on fetal and child development: A critical literature review. *Neuroscience & Biobehavioral Reviews, 43*, 137-162. doi:10.1016/j.neubiorev.2004.10.007
- Gröger, N., Matas, E., Gos, T., Lesse, A., Poeggel, G., Braun, K., & Bock, J. (2016). The transgenerational transmission of childhood adversity: behavioral, cellular, and epigenetic correlates. *Journal of Neural Transmission, 123*(9), 1037-1052. doi:10.1007/s00702-016-1570-1
- Hair, J. F., Black, W. C., Babin, B. J., Anderson, R. E., & Tatham, R. L. (2009). *Multivariate data analysis*. Upper Saddle River, N.J.: Pearson Education Limited.
- Hancock, G. R., & Mueller, R. O. (2001). Rethinking construct reliability within latent variable systems. In R. Cudeck, S. d. Toit, & D. Sörbom (Eds.), *Structural equation modeling: Present and future* (pp. 195-216). Lincolnwood, IL: Scientific Software International.

- Hayton, J. C., Allen, D. G., & Scarpello, V. (2004). Factor retention decisions in exploratory factor analysis: A tutorial on parallel analysis. *Organizational Research Methods, 7*(2), 191-205. doi:10.1177/1094428104263675
- Hernández-Martínez, C., Val, V. A., Murphy, M., Busquets, P. C., & Sans, J. C. (2011). Relation between positive and negative maternal emotional states and obstetrical outcomes. *Women & Health, 51*(2), 124-135. doi:10.1080/03630242.2010.550991
- Howard, L. M., Molyneaux, E., Dennis, C.-L., Rochat, T., Stein, A., & Milgrom, J. (2014). Non-psychotic mental disorders in the perinatal period. *The Lancet, 384*(9956), 1775-1788. doi:10.1016/S0140-6736(14)61276-9
- Howell, R. T., Kern, M. L., & Lyubomirsky, S. (2007). Health benefits: Meta-analytically determining the impact of well-being on objective health outcomes. *Health Psychology Review, 1*(1), 83-136. doi:10.1080/17437190701492486
- Hu, Y., Stewart-Brown, S., Twigg, L., & Weich, S. (2007). Can the 12-item General Health Questionnaire be used to measure positive mental health? *Psychological Medicine, 37*(7), 1005-1013. doi:10.1017/S0033291707009993
- Huizink, A. C., Mulder, E. J., & Buitelaar, J. K. (2004). Prenatal stress and risk for psychopathology: specific effects or induction of general susceptibility? *Psychological Bulletin, 130*(1), 115. doi:10.1037/0033-2909.130.1.115
- Huppert, F. A., & Whittington, J. E. (2003). Evidence for the independence of positive and negative well-being: Implications for quality of life assessment. *British Journal of Health Psychology, 8*(1), 107-122. doi:10.1348/135910703762879246
- Iwata, N., Mishima, N., Shimizu, T., Mizoue, T., Fukuhara, M., Hidano, T., & Spielberger, C. D. (1998). Positive and negative affect in the factor structure of the State-Trait Anxiety Inventory for Japanese workers. *Psychological Reports, 82*(2), 651-656. doi:10.2466/pr0.1998.82.2.651

- Karmaliani, R., Asad, N., Bann, C. M., Moss, N., McClure, E. M., Pasha, O., . . . Goldenberg, R. L. (2009). Prevalence of anxiety, depression and associated factors among pregnant women of hyderabad, pakistan. *International Journal of Social Psychiatry, 55*(5), 414-424. doi:10.1177/0020764008094645
- Kawabata, Y., Tseng, W.-L., & Crick, N. R. (2014). Mechanisms and processes of relational and physical victimization, depressive symptoms, and children's relational-interdependent self-construals: Implications for peer relationships and psychopathology. *Development and Psychopathology, 26*(3), 619-634. doi:10.1017/S0954579414000273
- Keyes, C. L. (2002). The mental health continuum: From languishing to flourishing in life. *Journal of Health and Social Behavior, 207-222*.
- Keyes, C. L., Dhingra, S. S., & Simoes, E. J. (2010). Change in level of positive mental health as a predictor of future risk of mental illness. *American Journal of Public Health, 100*(12), 2366-2371. doi:10.2105/AJPH.2010.192245
- Kvaal, K., Laake, K., & Engedal, K. (2001). Psychometric properties of the state part of the Spielberger State-Trait Anxiety Inventory (STAI) in geriatric patients. *International Journal of Geriatric Psychiatry, 16*(10), 980-986. doi:10.1002/gps.458
- Laake, L. M., & Bridgett, D. J. (2014). Happy babies, chatty toddlers: Infant positive affect facilitates early expressive, but not receptive language. *Infant Behavior and Development, 37*(1), 29-32. doi:10.1016/j.infbeh.2013.12.006
- Lamers, S. M., Westerhof, G. J., Glas, C. A., & Bohlmeijer, E. T. (2015). The bidirectional relation between positive mental health and psychopathology in a longitudinal representative panel study. *The Journal of Positive Psychology, 10*(6), 553-560. doi:10.1080/17439760.2015.1015156

- Le Strat, Y., Dubertret, C., & Le Foll, B. (2011). Prevalence and correlates of major depressive episode in pregnant and postpartum women in the United States. *Journal of Affective Disorders, 135*(1–3), 128-138. doi:10.1016/j.jad.2011.07.004
- McDougall, P., & Vaillancourt, T. (2015). Long-term adult outcomes of peer victimization in childhood and adolescence: Pathways to adjustment and maladjustment. *American Psychologist, 70*(4), 300. doi:10.1037/a0039174
- Moran, M. D. (2003). Arguments for rejecting the sequential bonferroni in ecological studies. *Oikos, 100*(2), 403-405.
- Moreno, A. J., & Robinson, J. L. (2005). Emotional vitality in infancy as a predictor of cognitive and language abilities in toddlerhood. *Infant and Child Development, 14*(4), 383-402. doi:10.1002/icd.406
- Mulder, E. J. H., Robles de Medina, P. G., Huizink, A. C., Van den Bergh, B. R. H., Buitelaar, J. K., & Visser, G. H. A. (2002). Prenatal maternal stress: effects on pregnancy and the (unborn) child. *Early Human Development, 70*(1–2), 3-14. doi:10.1016/S0378-3782(02)00075-0
- Muthén, L. K., & Muthén, B. O. (1998-2012). *Mplus User's Guide* (Seventh ed.). Los Angeles, CA: Muthén & Muthén.
- Newnham, E. A., Hooke, G. R., & Page, A. C. (2010). Progress monitoring and feedback in psychiatric care reduces depressive symptoms. *Journal of Affective Disorders, 127*(1), 139-146. doi:10.1016/j.jad.2010.05.003
- O'Donnell, K. J., & Meaney, M. J. (2017). Fetal Origins of Mental Health: The Developmental Origins of Health and Disease Hypothesis. *American Journal of Psychiatry, 174*(4), 319-328. doi:10.1176/appi.ajp.2016.16020138

- Parker, J. G., & Asher, S. R. (1987). Peer relations and later personal adjustment: Are low-accepted children at risk? *Psychological Bulletin*, *102*(3), 357. doi:10.1037/0033-2909.102.3.357
- Pearson, R. M., Evans, J., Kounali, D., Lewis, G., Heron, J., Ramchandani, P. G., . . . Stein, A. (2013). Maternal depression during pregnancy and the postnatal period: Risks and possible mechanisms for offspring depression at age 18 years. *Jama psychiatry*, *70*(12), 1312-1319. doi:10.1001/jamapsychiatry.2013.2163
- Pluess, M., Velders, F. P., Belsky, J., van IJzendoorn, M. H., Bakermans-Kranenburg, M. J., Jaddoe, V. W., . . . Tiemeier, H. (2011). Serotonin transporter polymorphism moderates effects of prenatal maternal anxiety on infant negative emotionality. *Biological Psychiatry*, *69*(6), 520-525. doi:10.1016/j.biopsych.2010.10.006
- Qiu, A., Tuan, T. A., Li, Y., Chen, H., Rifkin-Graboi, A., Broekman, B. F. P., . . . Meaney, M. J. (2015). Prenatal maternal depression alters amygdala functional connectivity in 6-month-old infants. *Translational psychiatry*, *5*, e508. doi:10.1038/tp.2015.3
- Qiu, A., Tuan, T. A., Ong, M. L., Li, Y., Chen, H., Rifkin-Graboi, A., . . . Meaney, M. J. (2015). COMT Haplotypes Modulate Associations of Antenatal Maternal Anxiety and Neonatal Cortical Morphology. *American Journal of Psychiatry*, *172*(2), 163-172. doi:10.1176/appi.ajp.2014.14030313
- Reise, S. P. (2012). The rediscovery of bifactor measurement models. *Multivariate Behavioral Research*, *47*(5), 667-696. doi:10.1080/00273171.2012.715555
- Rifkin-Graboi, A., Kong, L., Sim, L. W., Sanmugam, S., Broekman, B. F. P., Chen, H., . . . Qiu, A. (2015). Maternal sensitivity, infant limbic structure volume and functional connectivity: a preliminary study. *Translational psychiatry*, *5*, e668. doi:10.1038/tp.2015.133

- Rodriguez, A., Reise, S. P., & Haviland, M. G. (2016). Evaluating bifactor models: Calculating and interpreting statistical indices. *Psychological Methods, 21*(2), 137-150.
doi:10.1037/met0000045
- Ryff, C. D. (1989). Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *Journal of Personality and Social Psychology, 57*(6), 1069.
doi:10.1037/0022-3514.57.6.1069
- Santos, H., Tan, X., & Salomon, R. (2016). Heterogeneity in perinatal depression: how far have we come? A systematic review. *Archives of Women's Mental Health, 1*-13.
doi:10.1007/s00737-016-0691-8
- Schwartz, D., Lansford, J. E., Dodge, K. A., Pettit, G. S., & Bates, J. E. (2015). Peer victimization during middle childhood as a lead indicator of internalizing problems and diagnostic outcomes in late adolescence. *Journal of Clinical Child and Adolescent Psychology, 44*(3), 393-404. doi:10.1080/15374416.2014.881293
- Shafer, A. B. (2006). Meta-analysis of the factor structures of four depression questionnaires: Beck, CES-D, Hamilton, and Zung. *Journal of Clinical Psychology, 62*(1), 123-146.
doi:10.1002/jclp.20213
- Sherman, D. K., Bunyan, D. P., Creswell, J. D., & Jaremka, L. M. (2009). Psychological vulnerability and stress: The effects of self-affirmation on sympathetic nervous system responses to naturalistic stressors. *Health Psychology, 28*(5), 554. doi:10.1037/a0014663
- Simms, L. J., Grös, D. F., Watson, D., & O'hara, M. W. (2008). Parsing the general and specific components of depression and anxiety with bifactor modeling. *Depression and Anxiety, 25*(7), E34-E46. doi:10.1002/da.20432
- Soh, S. E., Lee, S. S. M., Hoon, S. W., Tan, M. Y., Goh, A., Lee, B. W., . . . Saw, S. M. (2012). The methodology of the GUSTO cohort study: A novel approach in studying pediatric allergy. *Asia Pacific Allergy, 2*(2), 144-148. doi:10.5415/apallergy.2012.2.2.144

- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the state-trait anxiety inventory*. Palo Alto, CA: Consulting Psychologists Press.
- Stephoe, A., Wright, C., Kunz-Ebrecht, S. R., & Iliffe, S. (2006). Dispositional optimism and health behaviour in community-dwelling older people: Associations with healthy ageing. *British Journal of Health Psychology, 11*(1), 71-84. doi:10.1348/135910705X42850
- Szekely, E., Pappa, I., Wilson, J. D., Bhamidi, S., Jaddoe, V. W., Verhulst, F. C., . . . Shaw, P. (2016). Childhood peer network characteristics: Genetic influences and links with early mental health trajectories. *Journal of Child Psychology and Psychiatry, 57*(6), 687-694. doi:10.1111/jcpp.12493
- Tam, H.-I., Yuk-ching, S. K. L., Hay-ming, H. L., Yiu-tsang, A. L., Yeung, W.-k., & Ip-ki, C. L. (2017). The moderating effects of positive psychological strengths on the relationship between parental anxiety and child depression: The significance of father's role in Hong Kong. *Children and Youth Services Review*. doi:10.1016/j.childyouth.2017.01.001
- Taylor, S. E., Lerner, J. S., Sherman, D. K., Sage, R. M., & McDowell, N. K. (2003). Are self-enhancing cognitions associated with healthy or unhealthy biological profiles? *Journal of Personality and Social Psychology, 85*(4), 605. doi:10.1037/0022-3514.85.4.605
- Van den Bergh, B. R. H., Mulder, E. J. H., Mennes, M., & Glover, V. (2005). Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: links and possible mechanisms. A review. *Neuroscience & Biobehavioral Reviews, 29*(2), 237-258. doi:10.1016/j.neubiorev.2004.10.007
- Waters, C. S., Hay, D. F., Simmonds, J. R., & van Goozen, S. H. (2014). Antenatal depression and children's developmental outcomes: Potential mechanisms and treatment options. *European Child and Adolescent Psychiatry, 23*(10), 957-971. doi:10.1007/s00787-014-0582-3

World Health Organization. (2004). *Promoting mental health: Concepts, emerging evidence, practice*. Retrieved from

<http://apps.who.int/iris/bitstream/10665/42940/1/9241591595.pdf>