

Integrating Metabolic Engineering and Heterogeneous Chemocatalysis: New Opportunities for Biomass to Chemicals**


Yanping Huo¹, Huaqiang Zeng^{2*}, Yugen Zhang^{2*}

((Dedication----optional))

Driven by the constant energy- and resource-saving demand, pursuing high step- and atom-economy is an endless process in chemical synthesis.¹ Recently, there are several comprehensive reviews involving the combination of biocatalysis and chemocatalysis.^{2,3} The combining of individual reaction steps into one-pot, even from the same type of technology such as chemocatalysis or biocatalysis, has great potential in terms of saving solvent, energy, time and space, as well as reducing waste production.^{3,4} The integration of reaction steps from different fields will hence be more attractive in unveiling new opportunities and achieving challenging tasks. Biocatalysis is catalyzed by enzymes and engineered metabolic processes which are typically carried out in water/media, at low concentration and has biogenic impurities from fermentation. Chemocatalysis with organo- or metal-catalysts is generally carried out in various solvents and catalysts which may sometimes be sensitive to impurities. One major challenge in combining biocatalysis and chemocatalysis lies in the compatibility of the two systems, including reaction conditions, conversions and selectivities.^{4,5} Although the groundbreaking pioneering work in this topic was reported by Bekkum group in 1980 illustrating a heterogenized metal catalyzed hydrogenation coupled with an enzymatic isomerization of D-glucose for the synthesis of D-mannitol, surprisingly this strategy has not been applied into biomass transformation till very recently.⁶

Biomass and derivatives generally are highly functionalized substrates with high oxygen content. Upgrading biomass to fuel and chemicals generally starts with reducing its oxygen content which by chemocatalysis often has less structure versatility.⁷ Alternatively, in biocatalysis especially metabolic process, biomass and its derivatives are served as substrates and/or nutrition in many natural processes. Various products with different carbon chain structures and different functionalities could be produced.⁸ The strategy of combining metabolic engineering and heterogeneous chemocatalysis is particularly suitable for biomass transformation process. With natural or engineered metabolic processes, biomass or derivatives can be selectively de-functionalized to platform intermediates that are suitable for chemocatalysis upgrading to value-added chemicals,

especially for those “drop-in” chemicals.



Scheme 1. Conversion of sugars to high-molecular-mass hydrocarbons by the combined metabolic engineering and chemocatalysis process.

Logically, there are different levels of combining/integrating biocatalysis and chemocatalysis that represent different levels of step- and atom-economy. The basic level corresponds to linking two individual processes with one key intermediate, which is generated from the first process but could serve as feed for the second reaction. This type of combining process is similar to the traditional cascade synthesis and the intermediate compound needs to be isolated and purified before being applied to the second step. Well-designed combination of metabolic process and chemocatalysis could be assembled together *via* engineering efforts to bridge the gap between two fields and achieve challenging tasks. Toste et al. reported an excellent example of this strategy for the conversion of sugars to jet fuel precursors in high yield.⁹ It is well known that short-chain alcohols, such as ethanol and *n*-butanol can be produced by natural biological routes at the yields approaching theoretical limit. However, to produce longer chain precursors of higher-molecular-mass hydrocarbons, which are commonly used in jet and diesel fuel, is still challenging to date. As shown in Scheme 1, *Clostridium acetobutylicum* was chosen by Toste et al. to produce acetone-*n*-butanol-ethanol (ABE) from sugars. The key is the combination of fermentation products where acetone has a nucleophilic α -carbon which is a suitable functionality for a metal catalyzed C-C bond formation with electrophilic alcohols present in ABE fermentation products. Glucose undergoes fermentation with *C. acetobutylicum* to give acetone, *n*-butanol and ethanol in 2.3/3.7/1 molar ratio at overall carbon yield of 90%. Fermentation products went through a two-step extraction and ABE was then passed through a Pd/C-K₃PO₄ heterogeneous catalyst system. Eventually, this combined system gave C₇-C₁₅ products in 38% overall carbon yield from glucose. The yield increased to 58% after inclusion of


[*] Dr. Y. Huo
Faculty of Chemical Engineering and Light Industry,
Guangdong University of Technology, Guangzhou,
Guangdong, 510006, China

Dr. H. Zeng, Y.G. Zhang
Institute of Bioengineering and Nanotechnology, 31
Biopolis Way, The Nanos, Singapore 138669, Singapore.
E-mail: hqzeng@ibn.a-star.edu.sg; yqzhang@ibn.a-star.edu.sg

[**] This work was supported by the Institute of Bioengineering and Nanotechnology, Biomedical Research Council, Agency for Science, Technology and Research.

C4-C6 products. Isolation and purification of fermentation products ABE is a critical issue in this combined system. It is energy intensive to extract ABE from water and to remove all impurities that may poison the metal catalyst.


Beckham et al. also demonstrated this strategy recently for the conversion of lignin to chemicals by combining biological and chemical catalysis (Scheme 2).^{10,11} Lignin is one of the major sources of biomass which is a heterogeneous polymer comprised of phenylpropanoid monomers. Due to its aromatics-rich heterogenized structure, lignin is very difficult to be selectively valorized. Chemically, lignin could be depolymerized by metal catalyzed reduction or oxidation under relatively harsh conditions.¹² However, some organisms use lignin-derived aromatic molecules as carbon sources in their metabolic pathways.¹³ Aromatic-catabolizing organisms convert pretreated lignin into central intermediates, such as catechol and protocatechuate. The dioxygenase enzymes then cleave C-C bonds in the aromatic ring to produce ring-opened species via β -ketoadipate pathway. Beckham et al.¹⁰ demonstrated this technology by using *Pseudomonas putida* to produce medium chain-length polyhydroxyalkanoates (*mcl*-PHAs) from lignin in an integrated biological process. *mcl*-PHA was extracted from cell culture system and re-precipitated from ethanol. The purified PHA underwent thermal depolymerization and then the depolymerized alkenoic acids were catalytically converted to hydrocarbons over a platinum-rhenium solid catalyst (300 °C, 27.5 bar H₂). In another work from the same group,¹¹ metabolic engineering of *P. putida* selectively produced muconate (molar yield 67%) from lignin-derived monomers. Muconic acid was isolated and purified from culture media through activated carbon absorption and recrystallization at low pH. Purified muconic acid then underwent metal-catalyzed hydrogenation to produce adipic acid (Pd/C, 24 bar H₂, 24 °C).¹⁴ These two examples opened a new window for upgrading lignin, a relatively inert biomass, by ring-opening of aromatic ring with biocatalysis and then further upgrading with chemocatalysis methods, although there are some difficulties in the connection between biocatalysis and chemocatalysis.



Scheme 2. The combining of metabolic engineering and chemocatalysis for lignin upgrading to hydrocarbons and adipic acid.

For this two-step biocatalysis-chemocatalysis process, one major drawback is that the intermediate substrates need to be isolated and purified to fulfill the requirement of the metal-catalyst system. This drawback will apparently affect the overall energy, mass and economic efficiency of the combined system. Dumesic et al. constantly worked on improving catalyst tolerance toward biogenic impurities by manipulating microenvironments of catalytically active sites of heterogeneous catalysts.¹⁵ Triacetic acid lactone (TAL) is a platform molecule derived from sugars by metabolic processes.¹⁶ In the process of further upgrading to sorbic acid via palladium or ruthenium catalyzed hydrogenation reactions, metal catalysts are very sensitive to the trace biogenic impurities

even after the TAL went through an intensive purification process. It was found that a poly(vinyl alcohol) (PVA) coating could create a microenvironment that is undesirable to biogenic impurities. Therefore, a PVA-overcoated metal catalyst was an order of magnitude more stable than catalyst without PVA coating in the hydrogenation of TAL. Efforts to minimize the gap between bio- and chemo-catalysis are definitely important. However, to develop new strategy allowing biocatalysis and chemocatalysis to work in one pot is more attractive and appreciable.



Scheme 3. The sequential fermentation/electrocatalytic reduction of glucose into 3-hexenedioic acid by a combined metabolic engineering and electrocatalysis process in one pot. The selected electrocatalysis step allows using fermentation broth as electrolyte and hydrogen source.

Due to the different nature of biocatalysis and chemocatalysis, combining two processes into one pot is very challenging. Recently, Tessonier et al.¹⁷ made a breakthrough in this field by introducing electro-chemocatalysis into this strategy and realized the combination of bio- and chemo-catalysis in one pot without any additional work-up in between. The process started from fermentation of glucose with engineered strain of *Saccharomyces cerevisiae* to muconic acid, a well-known metabolic process. High muconic acid titer of 559.5 mg/L was achieved with this new engineered strain. Muconic acid was then directly electrocatalytically hydrogenated to 3-hexenedioic acid (3-HAD) in a three-electrode electrochemical cell without any work up process. In this step, hydrogen as a reducing agent was produced *in situ* by water splitting on electrode, and the reaction occurred at ambient temperature and pressure. 94% yield of 3-HAD (overall yield of 67% from glucose) was obtained in the system with all the biogenic impurities. It was claimed that the charge on the electrode surface could help with the mitigation of poisoning. Although lead, a toxic metal, was used in this system to increase the resistance of electrode to biogenic impurities, the system did demonstrate an exciting potential in integrating biocatalysis and chemocatalysis in one pot for biomass upgrading to chemicals.

It is clear that there is great potential of applying integrated metabolic process and chemocatalysis systems in biomass transformation. Due to the great versatility of metabolic systems, different biomass resources could be converted into a variety of intermediate compounds. Chemical process can further turn these intermediates to various desired targets. It is an important step that this strategy moved from a two-step procedure to one pot reaction,

since it means tremendous savings in energy, mass, time and space. Further efforts may direct to develop real integrated system where biocatalysis and chemocatalysis work concurrently and synergistically in one system.

Received: ((will be filled in by the editorial staff))

Published online on ((will be filled in by the editorial staff))

Keywords: integrating system · metabolic engineering · chemocatalysis · biomass transformation

-
- [1] I. Arends, R. Sheldon, U. Hanfeld, *Green Chemistry and Catalysis*. Wiley-VCH; **2007**.
 - [2] A. Bruggink, R. Schoevaart, T. Kieboom, *Org. Proc. Res. Dev.* **2003**, 7, 622-640.
 - [3] a) O. Pamies, J. E. Bäckvall, *Chem. Rev.* **2003**, 103, 3247-3261; b) A. Corma, S. Iborra, A. Velty, *Chem. Rev.* **2007**, 107, 2411.
 - [4] a) H. Gröger, W. Hummel, *Curr. Opin. Chem. Biol.* **2014**, 19, 171; b) J. Nikolau, M. A. D. N. Perera, L. Brachova, B. Shanks, *Plant J.* **2008**, 54, 536.
 - [5] a) Z. Zhang, J. E. Jackson, D. J. Miller, *Bioresour. Technol.* **2008**, 99, 5873.
 - [6] a) M. Makkee, A. P. G. Kieboom, H. van Bekkum, J. A. Roels, *J C S Chem Commun* **1980**, 930-931; b) Y.-C. Lee, S. Dutta, K. C.-W. Wu, *ChemSusChem* **2014**, 7, 3241; c) H. Zhang, X. Li, X. Su, E. L. Ang, Y. Zhang, H. Zhao, *ChemCatChem*, **2016**, DOI: 10.1002/cctc.201600069.
 - [7] a) M. Besson, P. Gallezot and C. Pinel, *Chem. Rev.* **2014**, 114, 1827-1870; b) S. P. Teong, G. Yi, Y. Zhang, *Green Chem.* **2014**, 16, 2015-2026; c) A. Rahimi, A. Ulbrich, J. J. Coon, S. S. Stahl, *Nature*, **2014**, 515, 249-252; d) J. B. Binder, R. T. Raines, *Proc. Natl. Acad. Sci. USA* **2010**, 107, 4516-4521.
 - [8] J. D. Keasling, *Science* **2010**, 330, 1355-1358.
 - [9] P. Anbarasan, Z. C. Baer, S. Sreekumar, E. Gross, J. B. Binder, H. W. Blanch, D. S. Clark, F. D. Toste, *Nature* **2012**, 491, 235-239.
 - [10] J. G. Linger, D. R. Vardon, M. T. Guarnieri, E. M. Karp, G. B. Hunsinger, M. A. Franden, C. W. Johnson, G. Chupka, T. J. Strathmann, P. T. Pienkos, G. T. Beckham, *Proc. Natl. Acad. Sci. USA* **2014**, 111, 12013.
 - [11] D. R. Vardon, M. A. Franden, C. W. Johnson, E. M. Karp, M. T. Guarnieri, J. G. Linger, M. J. Salm, T. J. Strathmann, G. T. Beckham, *Energy Environ. Sci.* **2015**, 8, 617.
 - [12] J. Zakzeski, P. C. A. Bruijninx, A. L. Jongerius, B. M. Weckhuysen, *Chem. Rev.* **2010**, 110, 3552-3599.
 - [13] T. D. H. Bugg, M. Ahmad, E. M. Hardiman, R. Singh, *Curr. Opin. Biotechnol.* **2011**, 22, 394-400.
 - [14] a) M. Shiramizu, F. D. Toste, *Angew. Chem. Int. Ed.* **2013**, 52, 12905-12909; b) X. Li, D. Wu, G. Yi, H. Su, Y. Zhang, *Angew. Chem. Int. Ed.* **2014**, 53, 4200-4204.
 - [15] T. J. Schwartz, B. J. O'Neill, B. H. Shanks, J. A. Dumesic, *ACS Catal.* **2014**, 4, 2060.
 - [16] T. J. Schwartz, R. L. Johnson, J. Cardenas, A. Okerlund, N. A. Da Silva, K. Schmidt-Rohr, J. A. Dumesic, *Angew. Chem. Int. Ed.* **2014**, 53, 12718.
 - [17] M. Suastegui, J. E. Mathiesen, J. M. Carraher, N. Hernandez, N. R. Quiroz, A. Okerlund, E. W. Cochran, Z. Shao, J.-P. Tessonier, *Angew. Chem. Int. Ed.* **2016**, DOI: 10.1002/anie.201509653.

Entry for the Table of Contents (Please choose one layout)


Layout 2:

((Catch Phrase))

Yanping Huo ¹, Huaqiang Zeng^{2*},
Yugen Zhang^{*2}

_____ **Page – Page**

Integrating Metabolic Engineering and
Heterogeneous Chemocatalysis: New
Opportunities for Biomass to Chemicals



Integrating metabolic engineering process and chemocatalysis system creates tremendous opportunities in biomass transformation.