

# Shallow quantum circuits for efficient preparation of Slater determinants and correlated states on a quantum computer


Chong Hian Chee,<sup>1,\*</sup> Daniel Leykam,<sup>1</sup> Adrian M. Mak<sup>2</sup> and Dimitris G. Angelakis<sup>1,3,4,†</sup>

<sup>1</sup>Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543

<sup>2</sup>Institute of High Performance Computing, Agency for Science, Technology & Research (A\*STAR),  
1 Fusionopolis Way, 16-16 Connexis, Singapore 138632

<sup>3</sup>School of Electrical and Computer Engineering, Technical University of Crete, 73100 Chania, Greece

<sup>4</sup>AngelQ Quantum Computing, 531A Upper Cross Street, 04-95 Hong Lim Complex, Singapore 051531

 (Received 25 January 2023; revised 23 May 2023; accepted 27 July 2023; published xxxxxxxxx)

Fermionic *Ansatz* state preparation is a critical subroutine in many quantum algorithms such as the variational quantum eigensolver for quantum chemistry and condensed-matter applications. The shallowest circuit depth needed to prepare Slater determinants and correlated states to date scales at least linearly with respect to the system size  $N$ . Inspired by data-loading circuits developed for quantum machine learning, we propose an alternate paradigm that provides shallower, yet scalable,  $O(d \log_2^2 N)$  two-qubit gate-depth circuits to prepare such states with  $d$  fermions, offering a subexponential reduction in  $N$  over existing approaches in second quantization, enabling high-accuracy studies of  $d \ll O(N/\log_2^2 N)$  fermionic systems with larger basis sets on near-term quantum devices.

DOI: [10.1103/PhysRevA.00.002400](https://doi.org/10.1103/PhysRevA.00.002400)

## I. INTRODUCTION

Quantum computers promise the ability to solve hard many-body problems in quantum chemistry and condensed-matter physics, including the computation of ground-state energies and simulation of quantum dynamics [1–3]. The relevant quantum algorithms frequently involve quantum state preparation as a key step. For example, the success probability of quantum phase estimation is determined by the overlap of a trial *Ansatz* state with the eigenstate of interest [4–6]. Thus, efficient preparation of high-quality *Ansatz* states is crucial for many-body applications of quantum computing [7,8].

Most existing methods for preparing fermionic *Ansätze* use second quantization with Jordan-Wigner mapping [9] to efficiently represent the quantum many-body fermionic wavefunction using a number of qubits that scales linearly in the system size [1,2]. Widely-used fermionic *Ansätze* typically fall into two broad classes: The first class consists of hardware-efficient *Ansätze* which use parameterized hardware-native gates to minimize the depth of the quantum circuit [10], but are difficult to optimize [11] and do not guarantee an accurate representation of the desired quantum state [12]. The second class consists of problem-inspired *Ansätze* which are more promising and explicitly incorporate the physics of the system of interest, but require deeper circuits that scale polynomially in system size, which exacerbates errors due to quantum noise and decoherence [13–20], limiting state-of-the-art demonstrations to less than a hundred qubits [21,22], leaving studies of chemically-relevant molecular

systems requiring more than  $10^2$ – $10^3$  qubits well out of reach [23,24].

The shallowest general-purpose problem-inspired *Ansatz* states to date are mean-field Hartree-Fock states [13–15], which are Slater determinants that can be prepared using a mesh of fermionic single-excitation gates which have a linear  $O(N)$  two-qubit gate depth in the number of qubits  $N$ , as shown in Fig. 1(a). While Hartree-Fock states are efficiently simulatable using classical computers, they nevertheless serve as a useful starting point for quantum computers to prepare more interesting classically intractable correlated quantum *Ansätze*, such as the unitary coupled-cluster *Ansatz*, which incorporates quantum correlations by applying number-conserving multifermion excitation operators to a reference Hartree-Fock state [16–20,25].

Fermionic excitation operators are examples of Givens-rotation gates which perform rotations in a two-dimensional fermionic subspace of a larger Hilbert space and, together with their controlled variants, form a universal quantum gate set to realize any particle-conserving unitaries [26,27]. Therefore, such Givens-rotation gates have been helpful for preparing various fermionic states in quantum chemistry and condensed-matter applications [13–17,28–30]. Recently, such gates have also attracted interest in the context of quantum linear algebra, where they were used to construct shallow-depth “Clifford-loader” gates, which are linear combinations of anticommuting operators, as a means to efficiently encode  $d$ -dimensional subspaces of  $\mathbb{R}^N$  into an  $N$ -qubit state [31–34], enabling potential end-to-end quantum speedups for several quantum machine learning and linear-algebra problems, including determinant sampling and topological data analysis [31]. It is thus timely to consider whether the implementation of Clifford loaders via the Givens rotation is useful for preparing fermionic *Ansatz* states.

\*ch.chee@u.nus.edu

†dimitris.angelakis@gmail.com




FIG. 1. Different approaches for preparing  $d$ -occupied Slater determinants of  $N$  modes on a quantum computer, assuming Jordan-Wigner mapping. (a) Existing approaches use a linear-depth mesh of fermionic single-excitation gates to apply a fermionic basis transformation to a reference state [13–15]. (b) The proposed Clifford loaders via the Givens-rotation approach applies a sequence  $\hat{C}$  of gates  $d$  times to an all-zero state of  $N$  qubits  $|0\rangle^{\otimes N}$ . (c)  $\hat{C}$  consists of two products of multiple Givens rotations,  $\hat{D}$  and  $\hat{D}^\dagger$ , that sandwich a Pauli- $X$  gate on the first qubit. (d) The Givens rotations  $\hat{U}_{\mu\nu}$  are arranged in a binary tree. (e)  $\hat{U}_{\mu\nu}$  is decomposed using Pauli rotation gates  $R_x$  and  $R_z$  acting on qubit  $\mu$ , Hadamard gates  $H$  acting on qubit  $\nu$ , and CNOT ladders  $B_{\mu\nu}$  acting on all qubits between  $\mu$  and  $\nu$ .  $\theta$  and  $\vec{A}$  are scalar and vector parameters, respectively.

respect to the system size  $N$  for a sufficiently slowly growing  $d$ . We also show how this approach can be extended using the same preparation technique for Slater determinants to prepare fermionic  $L$ -wise correlated *Ansatz* states, which correlate between  $L$ -tuples of the fermionic modes in the *Ansatz*, to get quantum circuits that are shallower than that of the Slater determinant by at least a factor of  $L$ . Finally, to validate our correlated *Ansatz*, we shall demonstrate how the  $L = 2$  pairwise correlated *Ansatz* can be used to capture a significant fraction of correlation energy using an example of hydrogen chains up to  $N = 20$  qubits, for which pairwise electronic correlation is likely to be significant. Our results establish Clifford loaders constructed out of Givens rotations as a promising method for efficient, practical, and scalable preparation of fermionic *Ansatz* states for large quantum chemistry applications on near-term quantum computers.

## II. PREPARING SLATER DETERMINANTS USING SHALLOW CIRCUITS

We begin by showing how a Slater determinant with  $d$  occupied and  $N - d$  unoccupied fermionic modes can be prepared using a shallow quantum circuit. An arbitrary Slater determinant  $|\Psi_1\rangle$  is defined as [13,35–37]

$$|\Psi_1(A)\rangle := \prod_{l=1}^d \sum_{\mu=1}^N A_{\mu l} \hat{a}_\mu^\dagger |\text{vac}\rangle, \quad (1)$$

where  $A$  is an  $N \times d$  real matrix such that all  $d$  columns are orthogonal and normalized,  $|\text{vac}\rangle$  is a vacuum state, and  $\hat{a}_\mu^\dagger$  is a creation operator acting on the  $\mu$ th mode. While the definition in Eq. (1) is pedagogically convenient, it requires nonunitary operators  $\sum_{\mu=1}^N A_{\mu l} \hat{a}_\mu^\dagger$  which cannot be directly implemented on a quantum circuit. Consequently, the most efficient method to prepare Slater determinants to date is to perform a fermionic basis transformation to a reference Slater-determinant state for a given skew-Hermitian parameter matrix  $\kappa$  as

$$|\Psi_1(A)\rangle = \exp \left[ \sum_{\mu, \nu=1}^N \kappa_{\mu\nu} \hat{a}_\mu^\dagger \hat{a}_\nu \right] \prod_{r=1}^d a_r^\dagger |\text{vac}\rangle, \quad (2)$$

where the fermionic basis transformation is implemented as a linear-depth mesh of fermionic single-excitation gates  $\exp[\theta(\hat{a}_\mu^\dagger \hat{a}_\nu - \hat{a}_\nu^\dagger \hat{a}_\mu)]$ , as shown in Fig. 1(a) [14,15].

We propose to improve the circuit-depth efficiency of preparing Slater determinants using the equivalent form

$$|\Psi_1(A)\rangle := \prod_{l=1}^d \sum_{\mu=1}^N A_{\mu l} \hat{p}_\mu |\text{vac}\rangle, \quad (3)$$

as shown in Appendix A, where we use anticommuting operators

$$\hat{p}_\mu = \hat{a}_\mu^\dagger + \hat{a}_\mu, \quad (4)$$

with the relation  $\{\hat{p}_\mu, \hat{p}_\nu\} = 2\delta_{\mu\nu} \mathbf{I}$ , as shown in Appendix B. Using the anticommuting operators  $\hat{p}_\mu$  instead of  $\hat{a}_\mu^\dagger$  allows us to exploit the recent result in Ref. [31] that provides a shallow  $O(d \log_2 N)$  Givens-rotation gate-depth decomposition of the

133 Clifford loader

$$\hat{C}(\vec{A}_l) = \sum_{\mu=1}^N A_{\mu l} \hat{P}_\mu \quad (5)$$

134 for some normalized column  $\vec{A}_l$ . Applying the Clifford loaders  
 135  $\hat{C}(\vec{A}_l)$  in succession  $d$  times on a vacuum state  $|\text{vac}\rangle$ , each with  
 136 orthogonal columns  $\vec{A}_l, l = 1, \dots, d$ , from matrix  $A$  generates  
 137 the desired Slater determinant

$$|\Psi_1(A)\rangle = \prod_{l=1}^d \hat{C}(\vec{A}_l) |\text{vac}\rangle, \quad (6)$$

138 as shown in Fig. 1(b). In the case of  $d > \frac{N}{2}$ , we may apply the  
 139 Clifford loader  $N - d$  times instead, followed by occupation-  
 140 vacant mode swap to all Fock bases at the end, which is  
 141 equivalent to a Pauli-X bit flip to all qubits under the Jordan-  
 142 Wigner mapping. The Slater determinant in Eq. (6) can be  
 143 simplified to [31,38]

$$|\Psi_1(A)\rangle = \sum_{|B|=d} \det(A_B) |B\rangle, \quad (7)$$

144 where the sum is over all possible combinations of the ordered  
 145 set  $B$  containing  $d$  unique integers from 1 to  $N$ ,  $A_B$  is a  $d \times$   
 146  $d$  matrix minor of  $A$  whose row indexes are restricted by  $B$ ,  
 147 and  $|B\rangle$  denotes an  $N$ -mode Fock basis with occupied modes  
 148 indexed by  $B$ , as shown in Appendix C.

149 The Clifford loader  $\hat{C}(\vec{A}_l)$  in Eq. (5) can be decomposed as

$$\hat{C}(\vec{A}_l) = \hat{D}(\vec{A}_l) \hat{p}_1 \hat{D}^\dagger(\vec{A}_l), \quad (8)$$

150 as shown in Fig. 1(c), where the operator  $\hat{p}_1 = \hat{a}_1^\dagger + \hat{a}_1$  acting  
 151 on the first mode is sandwiched by two products of multiple  
 152 Givens rotations, termed elsewhere as unary data loaders  $\hat{D}$   
 153 [31,34,39]. Its conjugate transpose  $\hat{D}^\dagger$  is expressed as

$$\hat{D}^\dagger(\vec{A}_l) = \prod_{s=1}^{\lceil \log_2 N \rceil} \left[ \prod_{\mu, \nu \in \mathfrak{S}_s} \hat{U}_{\mu\nu}(\theta_{\mu\nu}^{(sl)}) \right], \quad (9)$$

154 where the Givens rotations

$$\hat{U}_{\mu\nu}(\theta) = \exp[\theta \hat{p}_\mu \hat{p}_\nu] \quad (10)$$

155 are arranged in a binary-tree pattern according to  
 156 the set of  $(\mu, \nu)$  indexes  $\mathfrak{S}_s = \{(\mu, \nu) | \mu = 2^s(k -$   
 157  $1) + 1, \nu = 2^{s-1}(2k - 1) + 1, k \in \mathbb{Z}^+ \setminus \{0\}\}$  for each sublayer  
 158  $s \in \{1, \dots, \lceil \log_2 N \rceil\}$ , as shown in Fig. 1(d).

159 By treating  $\vec{A}_l = (A_{1l}, \dots, A_{Nl})$  in Eq. (9) as a vector in the  
 160 basis of  $\{\hat{p}_1, \dots, \hat{p}_N\}$ , we exploit the Givens-rotation property  
 161 that

$$\hat{U}_{\mu\nu}(\theta) \hat{p}_r \hat{U}_{\mu\nu}^\dagger(\theta) = \begin{cases} \cos(2\theta) \hat{p}_r + \sin(2\theta) \hat{p}_\nu & r = \mu, \\ \cos(2\theta) \hat{p}_r - \sin(2\theta) \hat{p}_\mu & r = \nu, \\ \hat{p}_r & r \neq \mu, \nu, \end{cases} \quad (11)$$

162 to obtain the required rotation angles  $\theta_{\mu\nu}^{(sl)} = \frac{1}{2} \arctan \frac{A_{\nu l}^{(s)}}{A_{\mu l}^{(s)}}$  clas-  
 163 sically by numerically performing parallel Givens rotations on  
 164  $\vec{A}_l$  that correspond to the sequence in  $\hat{D}^\dagger$  that successively  
 165 zeros out the vector elements until the first element of  $\vec{A}_l$   
 166 becomes  $A_{1l} = 1$ , corresponding to  $\hat{p}_1$ .




FIG. 2. Linear-depth cascading CNOT ladder  $B$  in the decompo-  
 sition of the Givens-rotation gate  $\hat{U}$  is replaced by a nonequivalent  
 logarithmic-depth circuit that has no effect on  $\hat{U}$ .

Under the Jordan-Wigner mapping, the Givens rotation in  
 Eq. (10) maps to  $\exp[-i\theta \hat{Y}_\mu \hat{X}_\nu \otimes_{r=\mu+1}^{\nu-1} \hat{Z}_r]$ , which is a Pauli-  
 string rotation gate that can easily be implemented on a  
 quantum circuit [28,40], as shown in Fig. 1(e). The controlled-  
 NOT (CNOT) ladder  $B_{\mu\nu}$  in the gate decomposition of the  
 Givens rotation  $\hat{U}_{\mu\nu}$  serves the purpose of encoding the parity  
 of nonexciting qubits into the rotation gate, and it consists of  
 a cascade of CNOT gates [28,40]. However, this CNOT ladder  
 can be replaced by the nonequivalent binary-tree CNOT gate  
 arrangement [31] shown in Fig. 2, without any effect on the  
 Givens rotation  $\hat{U}_{\mu\nu}$ , thereby reducing the CNOT depth from  
 linear to logarithmic in  $N$ . Thus, by implementing  $d$  such  
 Clifford loaders with these Givens-rotation gates, we can pre-  
 pare Slater determinants with shallow  $O(d \log_2^2 N)$  two-qubit  
 gate-depth quantum circuits.

### III. EXTENSION TO A CORRELATED ANSATZ

Next, we extended the technique above by introducing a  
 method to incorporate  $L$ -wise correlations into the fermionic  
 Ansatz state preparation, where  $L = 1$  reduces to the Slater-  
 determinant case. The idea is to use a set of anticommuting  
 operators that contains non-particle-preserving multibody  
 Fock operators for the Clifford loaders in Eq. (5) and apply  
 the same technique undertaken in the Slater-determinant case.  
 Here, for simplicity we work with Pauli-string operators under  
 the Jordan-Wigner mapping. We extend the anticommuting  
 operator  $\hat{p}_\mu = \bigotimes_{r=1}^{\mu-1} \hat{Z}_r \hat{X}_\mu$  used to prepare the Slater deter-  
 minant previously to

$$\hat{p}_\mu^{(L)} = \bigotimes_{r=1}^{\mu-1} \hat{Z}_{rL} \bigotimes_{r'=L(\mu-1)+1}^{\mu L} \hat{X}_{r'} \quad (12)$$

to generate  $L$ -wise correlations. The modified operator  $\hat{p}_\mu^{(L)}$   
 has  $L$  Pauli-X terms and  $\mu$  Pauli-Z terms with modulo  $L$   
 indexes such that  $\hat{p}_\mu^{(L)}$  remains anticommuting  $\{\hat{p}_\mu^{(L)}, \hat{p}_\nu^{(L)}\} =$   
 $2\delta_{\mu\nu} \mathbf{I}$ , as shown in Appendix B. For example, the  $L = 2$   
 pairwise correlated anticommuting operator is given as  $\hat{p}_\mu^{(2)} =$   
 $\hat{Z}_2 \hat{Z}_4 \hat{Z}_6 \dots \hat{Z}_{2\mu-2} \hat{X}_{2\mu-1} \hat{X}_{2\mu}$ . Thus, we may prepare an  $N$ -mode  
 $d$ -occupied  $L$ -wise correlated state  $|\Psi_L\rangle$  by applying  $\frac{d}{L}$  Clif-  
 ford loaders  $\hat{C}_L$ ,

$$|\Psi_L(G)\rangle = \prod_{l=1}^{\frac{d}{L}} \hat{C}_L(\vec{G}_l) |\text{vac}\rangle, \quad (13)$$




FIG. 3. (a) A product of multiple Givens rotations  $\hat{D}_L^\dagger$  that is composed of Givens-rotation gates  $\hat{U}_{\mu,\nu}^{(L)}$  in a binary-tree arrangement on the quantum circuit. (b)  $\hat{U}_{\mu,\nu}^{(L)}$  is decomposed using Pauli rotation gates  $R_x$ , acting on qubit  $L\mu$ , and  $R_z$ , acting on qubit  $L(\mu-1)+1$ , and Hadamard gates  $H$  and CNOT ladders  $B_{\mu\nu}^{(L)}$  that act on all  $2L-1$  qubits in  $\{L(\mu-1)+1, \dots, L\mu-1\}$  and  $\{L(\nu-1)+1, \dots, L\nu\}$ , with  $B_{\mu\nu}^{(L)}$  also acting on  $\nu-\mu$  additional qubits in  $\{L\mu, L(\mu+1), L(\mu+2), \dots, L(\nu-1)\}$ .  $\theta$  and  $\vec{G}$  are scalar and vector parameters, respectively.

where  $G$  is an  $\frac{N}{L} \times \frac{d}{L}$  orthonormal matrix and  $\vec{G}_l$  is a column vector of  $G$ , which simplifies to

$$|\Psi_L(G)\rangle \simeq \sum_{|B'|=\frac{d}{L}} \det(G_{B'}) |B'_L\rangle \quad (14)$$

up to an unobserved global phase, where the sum is over all combinations of the ordered set  $B'$  containing  $\frac{d}{L}$  unique integers between 1 and  $\frac{N}{L}$ ,  $G_{B'}$  is a  $\frac{d}{L} \times \frac{d}{L}$  matrix minor of  $G$  whose rows are restricted to  $B'$ ,  $B'_L = \{L(j-1)+1, L(j-1)+2, \dots, Lj | j \in B'\}$ , and  $|B'_L\rangle$  denotes an  $N$ -mode Fock state with occupied modes indexed by  $B'_L$ , as shown in Appendix C.

The  $L$ -wise correlated Ansatz state  $|\Psi_L\rangle$  in Eq. (14) is similar to the Slater determinant  $|\Psi_1\rangle$  from Eq. (7) with regard to how the amplitudes are calculated but differs in the Fock states that have nonzero amplitudes. In the Slater-determinant case, all Fock states with particle number  $d$  will have nonzero amplitudes, while in the  $L=2$  pairwise case, all Fock states that have both particle number  $d$  and  $L=2$ -tuple neighboring occupations and neighboring vacancies will have nonzero amplitudes. For instance, in the case  $N=4$  and  $d=2$ , Fock states  $\{|0011\rangle, |1100\rangle\}$  will have nonzero amplitudes, and  $\{|0101\rangle, |0110\rangle, |1010\rangle, |1001\rangle\}$  will have zero amplitudes.

For a given normalized column  $\vec{G}_l$ , we may define a corresponding  $L$ -wise correlated Clifford loader  $\hat{C}_L(\vec{G}_l) = \hat{D}_L^\dagger(\vec{G}_l) \hat{p}_1^{(L)} \hat{D}_L(\vec{G}_l)$ , where  $\hat{p}_1^{(L)} = \bigotimes_{r=1}^L \hat{X}_r$  consists of  $L$  Pauli- $X$  gates that act on the first  $L$  qubits sandwiched by two products of multiple Givens rotations,  $\hat{D}_L$  and  $\hat{D}_L^\dagger$ . Each  $\hat{D}_L$  is composed of Givens-rotation gates  $\hat{U}_{\mu,\nu}^{(L)}(\theta) = \exp[\theta \hat{p}_{\mu\nu}^{(L)} \hat{p}_\nu^{(L)}]$  arranged in a binary-tree pattern similar to that shown in Fig. 3(a). Since the Givens-rotation gate  $\hat{U}_{\mu,\nu}^{(L)}(\theta)$  is a Pauli-string rotation gate, its gate decomposition [40] and the




FIG. 4. Estimated two-qubit gate depth per occupied mode  $d$  to prepare an  $N$ -mode Slater determinant  $|\Psi_1\rangle$  and an  $L=2$  pairwise correlated Ansatz state  $|\Psi_2\rangle$  using Clifford loaders compared to existing  $d$ -independent linear-depth approaches.

corresponding rotation angle can be obtained in a fashion similar to that in the Slater-determinant case, as shown Fig. 3(b). We refer readers to Appendix D for the explicit form of the Givens rotation in terms of Pauli operators.

#### IV. RESOURCE ANALYSIS

For simplicity, we treat all types of two-qubit gate depths as equal and assume there is no circuit compilation. We estimated that the overall two-qubit gate depth required to prepare an  $L$ -wise correlated Ansatz state  $|\Psi_L\rangle$  is  $\frac{2d}{L} \{ \lceil \log_2 \frac{N}{L} \rceil + (1+2 \log_2 L) \lceil \log_2 \frac{N}{L} \rceil \} \approx O(\frac{d}{L} \log_2^2 \frac{N}{L})$ , as shown in Appendix E. This shows that our approach to prepare an  $L$ -wise correlated Ansatz state  $|\Psi_L\rangle$  is shallower than a Slater determinant  $|\Psi_1\rangle$  by at least a factor of  $L$ , *ceteris paribus*.

We plotted the estimated two-qubit gate depth per occupied mode for Slater determinants  $|\Psi_1\rangle$  and pairwise correlated Ansatz states  $|\Psi_2\rangle$  on a quantum computer, as shown in Fig. 4, and compared the result to the previous linear-depth approach of preparing  $d=2, 16$ , and 128 occupied Slater determinants, which has a  $d$ -independent two-qubit gate depth of about  $2N$  [13–15]. Indeed, the crossover point  $d \approx \frac{N}{\log_2^2 N}$  shows that the minimum number of qubits  $N$  required to achieve a shallower circuit increases subexponentially with the number of occupied modes  $d$ . However, we highlight that this crossover point can be practically surpassed by near-term quantum devices such as superconducting qubits [41] and trapped ions [42] with fewer than  $10^5$  qubits for systems with  $2 \leq d \leq 128$  occupied fermionic modes, which is a sizable range that encompasses many systems of interest in quantum chemistry and condensed-matter physics. In general, our approach is suitable for problem classes that have a sufficiently slowly growing  $d \ll O(\frac{N}{\log_2^2 N})$ . One such problem is the computation of quantum observable quantities for fermionic systems for the complete-basis-set limit, where  $d$  is preserved but said quantities are computed for increasing values of  $N$  and extrapolated using various schemes to very large limits of  $N$  [43–45].

#### V. EXAMPLE: LINEAR HYDROGEN MOLECULAR CHAINS

To validate our  $L=2$  pairwise correlated Ansatz state, we numerically evaluated the fraction of the electronic correlation energy  $E_{\text{pair}}/E_{\text{corr}}$  captured by the optimized




FIG. 5. Numerically calculated fraction of the electronic correlation energy ratio  $E_{\text{pair}}/E_{\text{corr}}$  captured by an optimized pairwise correlated *Ansatz* state for hydrogen chains up to  $H_6$  at a fixed interatomic distance of 1.4 bohrs for various basis-set mixtures up to 20 qubits.

pairwise correlated *Ansatz* state  $|\Psi_2(G^*)\rangle$  for three linear hydrogen molecular chains ( $H_2$ ,  $H_4$ , and  $H_6$ ) with  $d = 2, 4$ , and 6 electrons at a fixed interatomic distance of 1.4 bohrs, where the pairwise electronic correlation is likely to be significant.  $E_{\text{pair}} = E_{\text{HF}} - \langle \hat{H}_e \rangle$  is the correlation energy beyond the mean-field energy  $E_{\text{HF}}$  of a molecule captured by  $|\Psi_2(G^*)\rangle$ , while  $E_{\text{corr}} = E_{\text{HF}} - E_{\text{FCI}}$  is the exact value of the correlation energy, where  $E_{\text{FCI}}$  is known as the full-configuration-interaction (FCI) energy.  $G^*$  is an optimized parameter matrix, obtained using a classical quasi-Newton Limited-memory Broyden-Fletcher-Goldfarb-Shanno algorithm with Bound constraints (L-BFGS-B) optimizer that minimizes the expectation of the electronic Hamiltonian  $\langle \hat{H}_e \rangle$ . We considered different mixtures of atomic basis sets [24], slater-type orbital approximated by 3 Gaussian primitives (STO-3G), split-valence (6-31G, 6-311G), correlation-consistent polarized valence double zeta (cc-pVDZ), augmented correlation-consistent polarized valence double zeta (aug-cc-pVDZ) for each hydrogen atom, resulting in system sizes ranging from 4 to 20 qubits. Such mixing of basis sets for each atom is a common strategy in computational quantum chemistry to reduce the resources required to achieve a desired precision [46]. All calculations were performed numerically using SCIPY [47], PYSCF [48], and PENNYLANE [49]. Figure 5 shows that a large fraction of the electronic correlation energy  $E_{\text{pair}}/E_{\text{corr}}$  is captured by the optimized pairwise correlated *Ansatz* state  $|\Psi_2(G^*)\rangle$ .

## VI. SUMMARY AND OUTLOOK

We proposed Givens-rotation-based Clifford loaders for efficient preparation of  $d$  occupied Slater determinants  $|\Psi_1\rangle$  of  $N$  modes using shallower  $O(d \log_2^2 N)$  two-qubit gate-depth quantum circuits. We also showed that by redefining new sets of the anticommutation operators  $\hat{p}_\mu$  for the Clifford loaders, the same technique can be used to prepare  $L$ -wise correlated *Ansatz* states  $|\Psi_L\rangle$  to yield quantum circuits shallower than those of Slater determinants by at least a factor of  $L$ . As demonstrated in the application of  $L = 2$  pairwise correlated *Ansatz* states to hydrogen chains,  $L$ -wise correlated states are potentially useful in fermionic systems with significant  $L$ -wise fermionic correlation, even though they are not expected to fully capture all the correlation energy. It will be interesting to generalize the Clifford loaders to other types of fermionic correlation while keeping the same shallow gate-depth scaling intact.

To the best of our knowledge, our approach to fermionic *Ansatz* state preparation offers a subexponential improvement in gate depth over existing methods in the second quantization with respect to system size  $N$  for fermionic problems where the number of occupied modes is  $d \ll O(\frac{N}{\log_2^2 N})$ . Nonetheless, our results have established Clifford loaders via Givens rotations as an efficient, yet practical and scalable, fermionic *Ansatz* state-preparation technique, which will enable the study of molecules and materials requiring larger basis-set sizes on near-term quantum devices.

## ACKNOWLEDGMENTS

This research is supported by the National Research Foundation, Singapore, and A\*STAR under its CQT Bridging Grant and Quantum Engineering Programme, Grant No. NRF2021-QEP2-02-P02, A\*STAR (No. 21709), and by the European Union's Horizon Programme (HORIZON-CL4-2021-DIGITALEMERGING-02-10), Grant Agreement No. 101080085, QCFD.

## APPENDIX A: PROOF OF EQUIVALENCE BETWEEN TWO DEFINITIONS OF AN ARBITRARY SLATER DETERMINANT

An arbitrary Slater determinant  $|\Psi_1(A)\rangle$  with  $d$  occupied and  $N - d$  unoccupied fermionic modes is defined as

$$|\Psi_1(A)\rangle = \prod_{l=1}^d \sum_{k=1}^N A_{lk} \hat{a}_k^\dagger |\text{vac}\rangle, \quad (\text{A1})$$

where  $A$  is an  $N \times d$  real matrix such that all the  $d$  columns are orthogonal and normalized,  $|\text{vac}\rangle$  is a vacuum state, and  $\hat{a}_k^\dagger$  is a creation operator acting on the  $k$ th mode. To begin, we expand the product in Eq. (A1) and replace the index  $k$  in the summation with indices  $k_1, \dots, k_d$ ,

$$\begin{aligned} |\Psi_1(A)\rangle &= \left( \sum_{k_d=1}^N A_{dk_d} \hat{a}_{k_d}^\dagger \right) \cdots \left( \sum_{k_1=1}^N A_{1k_1} \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (\text{A2}) \\ &= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \cdots A_{1k_1} \hat{a}_{k_d}^\dagger \cdots \hat{a}_{k_1}^\dagger |\text{vac}\rangle. \quad (\text{A3}) \end{aligned}$$

We want to show that Eq. (A3) is equivalent to the alternate definition of the Slater determinant,

$$|\Psi_{1,\text{alt}}(A)\rangle = \prod_{l=1}^d \sum_{k=1}^N A_{lk} (\hat{a}_k^\dagger + \hat{a}_k) |\text{vac}\rangle, \quad (\text{A4})$$

where we use an anticommuting operator  $\hat{a}_k^\dagger + \hat{a}_k$ . Starting from Eq. (A4), we expand the product and replace index  $k$  in the summation with indices  $k_1, \dots, k_d$ ,

$$\begin{aligned} |\Psi_{1,\text{alt}}(A)\rangle &= \left( \sum_{k_d=1}^N A_{dk_d} (\hat{a}_{k_d}^\dagger + \hat{a}_{k_d}) \right) \\ &\cdots \left( \sum_{k_1=1}^N A_{1k_1} (\hat{a}_{k_1}^\dagger + \hat{a}_{k_1}) \right) |\text{vac}\rangle. \quad (\text{A5}) \end{aligned}$$

347 Here, we consider evaluating the product of the two rightmost  
348 summation terms in Eq. (A5). We split the derivation into two  
349 cases,  $d = 1$  and  $d > 1$ . For  $d = 1$ , we note that  $\hat{a}_j|\text{vac}\rangle = 0$   
350 for any mode  $j$ ; thus, Eqs. (A3) and (A5) become trivially  
351 equivalent. For  $d > 1$  we have,

$$\left[ \sum_{k_2=1}^N A_{2k_2} (\hat{a}_{k_2}^\dagger + \hat{a}_{k_2}) \right] \left[ \sum_{k_1=1}^N A_{1k_1} (\hat{a}_{k_1}^\dagger + \hat{a}_{k_1}) \right]$$

$$= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} (\hat{a}_{k_2}^\dagger + \hat{a}_{k_2}) (\hat{a}_{k_1}^\dagger + \hat{a}_{k_1}) \quad (\text{A6})$$

$$= \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} (\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \hat{a}_{k_2}^\dagger \hat{a}_{k_1} + \hat{a}_{k_2} \hat{a}_{k_1}^\dagger + \hat{a}_{k_2} \hat{a}_{k_1}). \quad (\text{A7})$$

352 We then apply the fermionic commutation relation  $\{\hat{a}_\alpha, \hat{a}_\beta^\dagger\} =$   
353  $\delta_{\alpha\beta} \mathbf{I}$  to Eq. (A7) to get

$$\sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} (\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \delta_{k_2 k_1} \mathbf{I} + \hat{a}_{k_2} \hat{a}_{k_1}), \quad (\text{A8})$$

354 and since the columns of  $A$  are orthogonal, the inner product  
355 between any column  $i, j$  vanishes,  $\sum_k A_{ik} A_{jk} = 0$ , and we

thus have

$$\sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} (\hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger + \hat{a}_{k_2} \hat{a}_{k_1}). \quad (\text{A9})$$

Substituting Eq. (A9) back into Eq. (A5) and using  
 $\hat{a}_j|\text{vac}\rangle = 0$  for any mode  $j$  gives

$$|\Psi_{1,\text{alt}}(A)\rangle = \left( \sum_{k_d=1}^N A_{dk_d} (\hat{a}_{k_d}^\dagger + \hat{a}_{k_d}) \right)$$

$$\dots \left( \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle. \quad (\text{A10})$$

Henceforth, we consider the even- and odd- $d$  cases separately.  
First, assuming  $d$  is even, we can reapply the result in Eq. (A9)  
to the rest of the pairs of summation terms in Eq. (A10), which  
yields

$$|\Psi_{1,\text{alt}}(A)\rangle = \left( \sum_{k_d, k_{d-1}=1}^N A_{dk_d} A_{d-1k_{d-1}} \hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \right)$$

$$\dots \left( \sum_{k_2, k_1=1}^N A_{2k_2} A_{1k_1} \hat{a}_{k_2}^\dagger \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (\text{A11})$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \hat{a}_{k_d}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle. \quad (\text{A12})$$

Alternatively, if  $d$  is odd, we have

$$|\Psi_{1,\text{alt}}(A)\rangle = \left( \sum_{k_d=1}^N A_{dk_d} (\hat{a}_{k_d}^\dagger + \hat{a}_{k_d}) \right) \sum_{k_{d-1}, \dots, k_1=1}^N A_{d-1k_{d-1}} \dots A_{1k_1} \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle \quad (\text{A13})$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} (\hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger + \hat{a}_{k_d} \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger) |\text{vac}\rangle \quad (\text{A14})$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \left( \hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger + \delta_{k_d k_{d-1}} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger - \hat{a}_{k_{d-1}}^\dagger \hat{a}_{k_d} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (\text{A15})$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \left( \hat{a}_{k_d}^\dagger \hat{a}_{k_{d-1}}^\dagger \dots \hat{a}_{k_1}^\dagger - \hat{a}_{k_{d-1}}^\dagger \delta_{k_d k_{d-2}} \hat{a}_{k_{d-3}}^\dagger \dots \hat{a}_{k_1}^\dagger + \hat{a}_{k_{d-1}}^\dagger \hat{a}_{k_d} \hat{a}_{k_{d-2}}^\dagger \dots \hat{a}_{k_1}^\dagger \right) |\text{vac}\rangle \quad (\text{A16})$$

$$= \sum_{k_d, \dots, k_1=1}^N A_{dk_d} \dots A_{1k_1} \hat{a}_{k_d}^\dagger \dots \hat{a}_{k_1}^\dagger |\text{vac}\rangle, \quad (\text{A17})$$

where we have applied the fermionic commutation relation in Eq. (A14) and matrix orthogonality in Eqs. (A15) and (A16).  
Hence, by combining the even Eq. (A12) and odd Eq. (A17) results, we have established the equivalence between the alternative  
definition in Eq. (A4) and the original definition in Eq. (A1).

## APPENDIX B: PROOF OF ANTICOMMUTATION RELATIONS

Here, we shall show that  $\hat{p}_k = \hat{a}_k^\dagger + \hat{a}_k$  has the desired anticommutation relation  $\{\hat{p}_i, \hat{p}_j\} = 2\delta_{ij} \mathbf{I}$ :

$$\{\hat{p}_i, \hat{p}_j\} = \{\hat{a}_i^\dagger + \hat{a}_i, \hat{a}_j^\dagger + \hat{a}_j\} \quad (\text{B1})$$

$$= (\hat{a}_i^\dagger + \hat{a}_i)(\hat{a}_j^\dagger + \hat{a}_j) + (\hat{a}_j^\dagger + \hat{a}_j)(\hat{a}_i^\dagger + \hat{a}_i) \quad (\text{B2})$$

$$= \hat{a}_j^\dagger \hat{a}_j^\dagger + \hat{a}_i \hat{a}_j^\dagger + \hat{a}_i^\dagger \hat{a}_j + \hat{a}_j \hat{a}_i^\dagger + \hat{a}_j^\dagger \hat{a}_i + \hat{a}_i^\dagger \hat{a}_i + \hat{a}_j \hat{a}_j^\dagger \quad (\text{B3})$$

$$= \hat{a}_i \hat{a}_j^\dagger + \hat{a}_i^\dagger \hat{a}_j + \hat{a}_j \hat{a}_i^\dagger + \hat{a}_j^\dagger \hat{a}_i \quad (\text{B4})$$

$$= 2\delta_{ij} \mathbf{I} \quad (\text{shown}). \quad (\text{B5})$$

370 Next, we shall show that  $\hat{p}_\mu^{(L)} = \bigotimes_{r=1}^{\mu-1} \hat{Z}_{rL} \bigotimes_{r'=L(\mu-1)+1}^{\mu L} \hat{X}_{r'}$ , used to incorporate  $L$ -wise correlation into the Clifford loaders  
371 via the Givens-rotation approach, has the desired anticommutation relation  $\{\hat{p}_\mu^{(L)}, \hat{p}_\nu^{(L)}\} = 2\delta_{\mu\nu} \mathbf{I}$ . We shall split the derivation into  
372 two cases,  $\mu = \nu$  and  $\mu < \nu$ . Let us first consider  $\mu = \nu$ , where

$$2\hat{p}_\mu^{(L)} \hat{p}_\mu^{(L)} = 2 \left( \bigotimes_{r=1}^{\mu-1} \hat{Z}_{rL} \quad \bigotimes_{r'=L(\mu-1)+1}^{\mu L} \hat{X}_{r'} \right) \left( \bigotimes_{s=1}^{\mu-1} \hat{Z}_{sL} \quad \bigotimes_{s'=L(\mu-1)+1}^{\mu L} \hat{X}_{s'} \right) \quad (\text{B6})$$

$$= 2\mathbf{I} \quad (\text{shown}). \quad (\text{B7})$$

373 Second, without loss of generality, let us consider  $\mu < \nu$ , and we note that  $\hat{X}_\mu \hat{Z}_\mu = -i\hat{Y}_\mu$ :

$$\hat{p}_\mu^{(L)} \hat{p}_\nu^{(L)} = \left( \bigotimes_{r=1}^{\mu-1} \hat{Z}_{rL} \quad \bigotimes_{r'=L(\mu-1)+1}^{\mu L} \hat{X}_{r'} \right) \left( \bigotimes_{s=1}^{\nu-1} \hat{Z}_{sL} \quad \bigotimes_{s'=L(\nu-1)+1}^{\nu L} \hat{X}_{s'} \right) \quad (\text{B8})$$

$$= -i \left[ \bigotimes_{r=(\mu-1)L+1}^{\mu L-1} \hat{X}_r \hat{Y}_{\mu L} \quad \bigotimes_{r'=\mu+1}^{\nu-1} \hat{Z}_{r'L} \quad \bigotimes_{r''=(\nu-1)L+1}^{\nu L} \hat{X}_{r''} \right] \quad (\text{B9})$$

374 Therefore, by noting  $\hat{Z}_\mu \hat{X}_\mu = i\hat{Y}_\mu$  we then have

$$\{\hat{p}_\mu^{(L)}, \hat{p}_\nu^{(L)}\} = \hat{p}_\mu^{(L)} \hat{p}_\nu^{(L)} + \hat{p}_\nu^{(L)} \hat{p}_\mu^{(L)} \quad (\text{B10})$$

$$= -i \left[ \bigotimes_{r=(\mu-1)L+1}^{\mu L-1} \hat{X}_r \hat{Y}_{\mu L} \quad \bigotimes_{r'=\mu+1}^{\nu-1} \hat{Z}_{r'L} \quad \bigotimes_{r''=(\nu-1)L+1}^{\nu L} \hat{X}_{r''} \right] \\ + i \left[ \bigotimes_{s=(\mu-1)L+1}^{\mu L-1} \hat{X}_s \hat{Y}_{\mu L} \quad \bigotimes_{s'=\mu+1}^{\nu-1} \hat{Z}_{s'L} \quad \bigotimes_{s''=(\nu-1)L+1}^{\nu L} \hat{X}_{s''} \right] \quad (\text{B11})$$

$$= 0 \quad (\text{shown}). \quad (\text{B12})$$

### APPENDIX C: EVALUATION OF THE SLATER DETERMINANT AND $L$ -WISE CORRELATED ANSATZ USING CLIFFORD LOADERS

375 In the main text, we proposed to construct Clifford loaders  
376  $\hat{C}$  using Givens rotation to prepare the Slater determinant  
377  $|\Psi_1(A)\rangle$  as follows:  
378

$$|\Psi_1(A)\rangle = \prod_{l=1}^d \hat{C}(\vec{A}_l) |\text{vac}\rangle. \quad (\text{C1})$$

381 We want to show that Eq. (C1) can be mathematically eval-  
382 uated using geometric algebra, also known as real Clifford  
383 algebra. We define a Clifford loader  $\hat{C}(\vec{x})$  for a given normal-  
384 ized size- $N$  vector  $\vec{x}$  as a linear combination of anticommuting  
385 operators  $\hat{p}_r$ , as follows:

$$\hat{C}(\vec{x}) = \sum_{\mu=1}^N x_\mu \hat{p}_\mu. \quad (\text{C2})$$

386 A geometric product of two Clifford loaders  $\hat{C}(\vec{x})\hat{C}(\vec{y})$  for  
387 any two normalized size- $N$  vectors  $\vec{x}$  and  $\vec{y}$  is defined as

$$\hat{C}(\vec{x})\hat{C}(\vec{y}) = \hat{C}(\vec{x}) \cdot \hat{C}(\vec{y}) + \hat{C}(\vec{x}) \wedge \hat{C}(\vec{y}), \quad (\text{C3})$$

388 where  $\cdot$  and  $\wedge$  refer to the standard inner dot and exterior  
389 wedge products, respectively. Substituting the definition in  
390 Eq. (C2) into Eq. (C3), we have

$$\hat{C}(\vec{x})\hat{C}(\vec{y}) = \sum_{r=1}^N x_r y_r (\hat{p}_r \cdot \hat{p}_r) + \sum_{\mu, \nu=1}^N x_\mu y_\nu (\hat{p}_\mu \wedge \hat{p}_\nu). \quad (\text{C4})$$

391 In this work, we shall consider only orthogonal and nor-  
392 malized size- $N$  vectors  $\vec{A}_l$  for  $l = 1, 2, \dots, d$ , such that the  
393 inner product of any two vectors is zero. As a result, the geo-  
394 metric product of two Clifford loaders is simply equivalent to  
395 its exterior product as the first term of Eq. (C4) vanishes under  
396 orthogonality. Thus, products of multiple Clifford loaders can  
397 be easily written as exterior products of multiple anticommut-  
398 ing operators,

$$\prod_{l=1}^d \hat{C}(\vec{A}_l) = \sum_{\mu, \nu, \dots, r=1}^N A_{\mu 1} A_{\nu 2} \cdots A_{rd} \underbrace{(\hat{p}_\mu \wedge \hat{p}_\nu \wedge \cdots \wedge \hat{p}_r)}_{d \text{ operators}}. \quad (\text{C5})$$

We note the following identities of the exterior product:

$$\hat{p}_\mu \wedge \hat{p}_\mu = 0 \quad (\text{C6})$$

400 and

$$\hat{p}_{\sigma_1} \wedge \hat{p}_{\sigma_2} \wedge \cdots \wedge \hat{p}_{\sigma_d} = \text{sgn}(\sigma) \hat{p}_{B_1} \wedge \hat{p}_{B_2} \wedge \cdots \wedge \hat{p}_{B_d}, \quad (\text{C7})$$

401 where we let  $\sigma$  be a permutation of  $\{B_1, B_2, \dots, B_d\}$  for any  
402 ordered set  $B$  containing  $d$  unique integers between 1 and  $N$   
and  $B_\mu$  and  $\sigma_\mu$  refer to the  $\mu$ th integers of  $B$  and  $\sigma$ , respectively. Using identities (C6) and (C7), the sum in Eq. (C5) reduces to

$$\prod_{l=1}^d \hat{C}(\vec{A}_l) = \sum_{|B|=d} \sum_{\sigma \in B} \text{sgn}(\sigma) A_{\sigma_1 1} A_{\sigma_2 2} \cdots A_{\sigma_d d} \times (\hat{p}_{B_1} \wedge \hat{p}_{B_2} \wedge \cdots \wedge \hat{p}_{B_d}), \quad (\text{C8})$$

where the outer sum is over all possible combinations of the ordered set  $B$  containing  $d$  unique integers between 1 and  $N$  and the inner sum is over all possible integer permutations  $\sigma$  of each  $B$ . Using the Leibniz-determinant formula for matrix minors

$$\det(A_B) = \sum_{\sigma \in B} \text{sgn}(\sigma) A_{\sigma_1 1} A_{\sigma_2 2} \cdots A_{\sigma_d d}, \quad (\text{C9})$$

where  $A_B$  is a  $d \times d$  matrix minor of  $A$  whose rows are restricted to  $B$ , we have

$$\prod_{l=1}^d \hat{C}(\vec{A}_l) = \sum_{|B|=d} \det(A_B) \hat{p}_{B_1} \wedge \hat{p}_{B_2} \wedge \cdots \wedge \hat{p}_{B_d}. \quad (\text{C10})$$

$$\hat{p}_\mu^{(L)} = \begin{cases} \bigotimes_{r=1}^{\mu-1} (\mathbf{I} - 2\hat{a}_{rL}^\dagger \hat{a}_{rL}) \bigotimes_{r'=L(\mu-1)+1}^{\mu L} [\hat{a}_{r'}^\dagger + (-1)^{r'} \hat{a}_{r'}] & \text{if } L \text{ is even,} \\ -\bigotimes_{r=1 \notin L\mathbb{Z}}^{L(\mu-1)-1} (\mathbf{I} - 2\hat{a}_r^\dagger \hat{a}_r) \bigotimes_{r'=L(\mu-1)+1}^{\mu L} [\hat{a}_{r'}^\dagger + (-1)^{r'-L(\mu-1)-1} \hat{a}_{r'}] & \text{if } L \text{ is odd,} \end{cases} \quad (\text{C14})$$

434 where the  $r$  index in the odd- $L$  case increments with size 1  
435 from 1 to  $L(\mu-1)-1$  but skips every index that is a multiple  
436 of  $L$ . Here, we note that in Eq. (C14), all fermionic terms in  
437 normal ordered form that contain any annihilation operators  
438 will vanish when acted upon by a vacuum state and fermionic  
439 terms that contain only creation operators will survive. Thus,  
440 by substituting Eq. (C14) into the product of  $\frac{d}{L}$  Clifford loaders  
441 in Eq. (C10) and applying it to a vacuum state  $|\text{vac}\rangle$ , we  
442 obtain an expression of the  $L$ -wise correlated *Ansatz* state up  
443 to an unobservable global phase,

$$\prod_{l=1}^{\frac{d}{L}} \hat{C}_L(\vec{G}_l) |\text{vac}\rangle \simeq \sum_{|B'|=\frac{d}{L}} \det(G_{B'}) |B'_L\rangle, \quad (\text{C15})$$

444 where the sum is over all possible combinations of ordered  
445 set  $B'$  containing  $\frac{d}{L}$  unique integers between 1 and  $\frac{N}{L}$ ,  $G_{B'}$  is  
446 a  $\frac{d}{L} \times \frac{d}{L}$  matrix minor of  $G$  whose rows are restricted to  $B'$ ,  
447  $B'_L = \{L(j-1)+1, L(j-1)+2, \dots, Lj | j \in B'\}$ , and  $|B'_L\rangle$   
448 denotes an  $N$ -mode Fock basis whose occupied modes are  
449 indexed by  $B'_L$ .

Hence, by letting the anticommuting operator be

$$\hat{p}_\mu = \hat{a}_\mu^\dagger + \hat{a}_\mu \quad (\text{C11})$$

and applying the product of  $d$  Clifford loaders (C10) to a vacuum state  $|\text{vac}\rangle$ , we obtain the alternative expression of the Slater determinant,

$$\prod_{l=1}^d \hat{C}(\vec{A}_l) |\text{vac}\rangle = \sum_{|B|=d} \det(A_B) |B\rangle, \quad (\text{C12})$$

where  $|B\rangle$  denotes a Fock state whose occupied modes are indexed by  $B$ .

Next, we consider extending the application of Clifford loaders to prepare  $L$ -wise correlated *Ansatz* states, where  $L=1$  reduces to the Slater-determinant case. The idea is to use a new set of anticommuting operators  $\hat{p}_\mu^{(L)}$  that contains non-particle-preserving multibody Fock operators for the Clifford loaders in Eq. (C2). In the main text, under the Jordan-Wigner mapping, we modify the anticommuting operator  $\hat{p}_\mu = \bigotimes_{r=1}^{\mu-1} \hat{Z}_r \hat{X}_\mu$  in Eq. (C11) used to prepare the Slater determinant above to

$$\hat{p}_\mu^{(L)} = \bigotimes_{r=1}^{\mu-1} \hat{Z}_{rL} \bigotimes_{r'=L(\mu-1)+1}^{\mu L} \hat{X}_{r'} \quad (\text{C13})$$

to apply  $L$ -wise correlation. This modified operator  $\hat{p}_\mu^{(L)}$  has  $L$  Pauli- $X$  terms and  $\mu$  Pauli- $Z$  terms with modulo  $L$  indexes such that  $\{\hat{p}_\mu^{(L)}, \hat{p}_\nu^{(L)}\} = 2\delta_{\mu\nu} \mathbf{I}$ . For example, the  $L=2$  pairwise correlated anticommuting operators are given as  $\hat{p}_\mu^{(2)} = \hat{Z}_2 \hat{Z}_4 \hat{Z}_6 \cdots \hat{Z}_{2\mu-2} \hat{X}_{2\mu-1} \hat{X}_{2\mu}$ . In terms of Fermionic creation and annihilation operators, Eq. (C13) maps back to

#### APPENDIX D: GIVENS-ROTATION GATE AND ITS DECOMPOSITION FOR THE $L$ -WISE CORRELATED ANSATZ STATE

The Givens-rotation gate is defined in the main text as

$$\hat{U}_{\mu\nu}^{(L)}(\theta) = \exp[\theta \hat{p}_\mu^{(L)} \hat{p}_\nu^{(L)}], \quad (\text{D1})$$

where, with the use of anticommuting operators in Eq. (C13), it becomes

$$\hat{U}_{\mu\nu}^{(L)}(\theta) = \exp \left[ -i\theta \bigotimes_{r=L(\mu-1)+1}^{\mu L-1} \hat{X}_r \hat{Y}_{\mu L} \bigotimes_{r'=L(\nu-1)+1}^{\nu L} \hat{X}_{r'} \bigotimes_{r''=\mu+1}^{\nu-1} \hat{Z}_{r''L} \right]. \quad (\text{D2})$$

For example, an  $L=2$  pairwise Givens rotation would be

$$\hat{U}_{\mu\nu}^{(2)}(\theta) = \exp[-i\theta \hat{X}_{2\mu-1} \hat{Y}_{2\mu} \hat{X}_{2\nu-1} \hat{X}_{2\nu} \hat{Z}_{2(\mu+1)} \times \hat{Z}_{2(\mu+2)} \cdots \hat{Z}_{2(\nu-1)}], \quad (\text{D3})$$

which can easily be decomposed, as shown in Fig. 6, using gate-decomposition techniques from [40]. Therefore, the






FIG. 6. The pairwise Givens-rotation gate  $\hat{U}_{\mu\nu}^{(2)}$  is decomposed using Pauli-rotation gates  $R_x$ , which acts on qubit  $2\mu$  that correspond to  $\hat{Y}_{2\mu}$ , and  $R_z$ , which acts on qubit  $2\mu - 1$ , and Hadamard gates  $H$  and CNOT ladders  $B_{\mu\nu}^{(L)}$  that act on all three qubits in  $\{2\mu - 1, 2\nu - 1, 2\nu\}$  that correspond to  $\hat{X}_{2\mu-1}\hat{X}_{2\nu-1}\hat{X}_{2\nu}$ , with  $B_{\mu\nu}^{(L)}$  also acting on  $\nu - \mu$  additional qubits in  $\{2\mu, 2(\mu+1), 2(\mu+2), \dots, 2(\nu-1)\}$ .  $\theta$  is a scalar parameter.

459 Givens-rotation gate  $\hat{U}_{\mu\nu}^{(L)}(\theta)$  used in this work is simply a  
460 Pauli-string-rotation gate whose gate decomposition is a gen-  
461 eralization of Fig. 6, as shown in Fig. 3(b) of the main text.

#### 462 APPENDIX E: DERIVATION OF THE TWO-QUBIT 463 GATE-DEPTH SCALING OF THE $L$ -WISE CORRELATED 464 QUANTUM CIRCUIT ANSATZ

465 To prepare the  $L$ -wise correlated *Ansatz* state  $|\Psi_L\rangle$  on a  
466 quantum computer under Jordan-Wigner mapping, we apply  
467  $\frac{d}{L}$  Clifford loaders  $\hat{C}_L$  to an all-zero qubit state. Each Clifford  
468 loader  $\hat{C}$  has two products of multiple Givens rotations,  $D_L$   
469 and  $D_L^\dagger$ . Each  $D_L$  has  $\lceil \log_2 \frac{N}{L} \rceil$  Givens-rotation gate depth.  
470 Each Givens-rotation gate  $U_{\mu\nu}^{(L)}$  contains two CNOT ladders

$B_{\mu\nu}^{(L)}$  that act on all  $2L + \nu - \mu - 1$  qubits in  $\{L(\mu - 1) +$   
471  $1, \dots, L\mu\}$ ,  $\{L(\mu + 1), L(\mu + 2), \dots, L(\nu - 1)\}$ , and  $\{L(\nu -$   
472  $1) + 1, \dots, L\nu\}$ . We employ the logarithmic-depth CNOT lad-  
473 ders shown in Fig. 2 in the main text; thus, each CNOT ladder  
474  $B_{\mu\nu}^{(L)}$  has a two-qubit gate depth of  $\lceil \log_2(2L + \nu - \mu - 1) \rceil \approx$   
475  $O(\log_2 N)$ . Focusing on all Givens rotations  $U_{1,2^s}^{(L)}$  that act  
476 on the first qubit in every sublayer  $s \in \{1, 2, \dots, \lceil \log_2 \frac{N}{L} \rceil\}$ , as  
477 shown in Fig. 3(a) in the main text, the overall two-qubit gate  
478 depth of the quantum circuit required to prepare the  $L$ -wise  
479 correlated *Ansatz* state  $|\Psi_L\rangle$  is estimated to be  
480

$$\frac{2d}{L} \sum_{s=1}^{\lceil \log_2 \frac{N}{L} \rceil} 2 \lceil \log_2(2L + 2^s - 1) \rceil$$

$$\leq \frac{4d}{L} \sum_{s=1}^{\lceil \log_2 \frac{N}{L} \rceil} \log_2(2^{s+\log_2(L)}) \quad (E1)$$

$$= \frac{4d}{L} \sum_{s=1}^{\lceil \log_2 \frac{N}{L} \rceil} [s + \log_2(L)] \quad (E2)$$

$$= \frac{2d}{L} \left( \left\lceil \log_2 \frac{N}{L} \right\rceil + (1 + 2 \log_2 L) \left\lceil \log_2 \frac{N}{L} \right\rceil \right). \quad (E3)$$

- [1] B. Bauer, S. Bravyi, M. Motta, and G. K.-L. Chan, Quantum algorithms for quantum chemistry and quantum materials science, *Chem. Rev.* **120**, 12685 (2020).
- [2] M. Motta and J. E. Rice, Emerging quantum computing algorithms for quantum chemistry, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **12**, e1580 (2022).
- [3] A. J. Daley, I. Bloch, C. Kokail, S. Flannigan, N. Pearson, M. Troyer, and P. Zoller, Practical quantum advantage in quantum simulation, *Nature (London)* **607**, 667 (2022).
- [4] D. S. Abrams and S. Lloyd, Simulation of Many-Body Fermi Systems on a Universal Quantum Computer, *Phys. Rev. Lett.* **79**, 2586 (1997).
- [5] D. S. Abrams and S. Lloyd, Quantum Algorithm Providing Exponential Speed Increase for Finding Eigenvalues and Eigenvectors, *Phys. Rev. Lett.* **83**, 5162 (1999).
- [6] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, Simulated quantum computation of molecular energies, *Science* **309**, 1704 (2005).
- [7] S. Lee, J. Lee, H. Zhai, Y. Tong, A. M. Dalzell, A. Kumar, P. Helms, J. Gray, Z.-H. Cui, W. Liu, M. Kastoryano, R. Babbush, J. Preskill, D. R. Reichman, E. T. Campbell, E. F. Valeev, L. Lin, and G. K.-L. Chan, Evaluating the evidence for exponential quantum advantage in ground-state quantum chemistry *Nat. Commun.* **14**, 1952 (2023).
- [8] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, W.-K. Mok, S. Sim, L.-C. Kwek, and A. Aspuru-Guzik, Noisy intermediate-scale quantum algorithms, *Rev. Mod. Phys.* **94**, 015004 (2022).
- [9] P. Jordan and E. Wigner, Über das Paulische Äquivalenzverbot, *Z. Phys.* **47**, 631 (1928).
- [10] A. Kandala, A. Mezzacapo, K. Temme, M. Takita, M. Brink, J. M. Chow, and J. M. Gambetta, Hardware-efficient variational quantum eigensolver for small molecules and quantum magnets, *Nature (London)* **549**, 242 (2017).
- [11] L. Bittel and M. Kliesch, Training Variational Quantum Algorithms is NP-Hard, *Phys. Rev. Lett.* **127**, 120502 (2021).
- [12] J. Tilly, H. Chen, S. Cao, D. Picozzi, K. Setia, Y. Li, E. Grant, L. Wossnig, I. Rungger, G. H. Booth, and J. Tennyson, The variational quantum eigensolver: A review of methods and best practices, *Phys. Rep.* **986**, 1 (2022).
- [13] D. Wecker, M. B. Hastings, N. Wiebe, B. K. Clark, C. Nayak, and M. Troyer, Solving strongly correlated electron models on a quantum computer, *Phys. Rev. A* **92**, 062318 (2015).
- [14] I. D. Kivlichan, J. McClean, N. Wiebe, C. Gidney, A. Aspuru-Guzik, G. K.-L. Chan, and R. Babbush, Quantum Simulation of Electronic Structure with Linear Depth and Connectivity, *Phys. Rev. Lett.* **120**, 110501 (2018).
- [15] F. Arute *et al.*, Hartree-Fock on a superconducting qubit quantum computer, *Science* **369**, 1084 (2020).
- [16] A. Anand, P. Schleich, S. Alperin-Lea, P. W. K. Jensen, S. Sim, M. Díaz-Tinoco, J. S. Kottmann, M. Degroote, A. F. Izmaylov, and A. Aspuru-Guzik, A quantum computing view on unitary coupled cluster theory, *Chem. Soc. Rev.* **51**, 1659 (2022).
- [17] F. A. Evangelista, G. K.-L. Chan, and G. E. Scuseria, Exact parameterization of fermionic wave functions via unitary coupled cluster theory, *J. Chem. Phys.* **151**, 244112 (2019).

- [18] Q. Wang, M. Li, C. Monroe, and Y. Nam, Resource-optimized fermionic local-Hamiltonian simulation on a quantum computer for quantum chemistry, *Quantum* **5**, 509 (2021).
- [19] J. S. Kottmann and A. Aspuru-Guzik, Optimized low-depth quantum circuits for molecular electronic structure using a separable-pair approximation, *Phys. Rev. A* **105**, 032449 (2022).
- [20] H. L. Tang, V. O. Shkolnikov, G. S. Barron, H. R. Grimsley, N. J. Mayhall, E. Barnes, and S. E. Economou, Qubit-ADAPT-VQE: An Adaptive Algorithm for Constructing Hardware-Efficient Ansätze on a Quantum Processor, *PRX Quantum* **2**, 020310 (2021).
- [21] T. E. O'Brien *et al.*, Purification-based quantum error mitigation of pair-correlated electron simulations, [arXiv:2210.10799](https://arxiv.org/abs/2210.10799).
- [22] R. N. Tazhigulov, S.-N. Sun, R. Haghshenas, H. Zhai, A. T. Tan, N. C. Rubin, R. Babbush, A. J. Minnich, and G. K.-L. Chan, Simulating Models of Challenging Correlated Molecules and Materials on the Sycamore Quantum Processor, *PRX Quantum* **3**, 040318 (2022).
- [23] V. E. Elfving, B. W. Broer, M. Webber, J. Gavartin, M. D. Halls, K. P. Lorton, and A. Bochevarov, How will quantum computers provide an industrially relevant computational advantage in quantum chemistry? [arXiv:2009.12472](https://arxiv.org/abs/2009.12472).
- [24] B. Nagy and F. Jensen, in *Reviews in Computational Chemistry* (John Wiley & Sons, Ltd., 2017), Vol. 30, pp. 93–149.
- [25] J. Romero, R. Babbush, J. R. McClean, C. Hempel, P. J. Love, and A. Aspuru-Guzik, Strategies for quantum computing molecular energies using the unitary coupled cluster ansatz, *Quantum Sci. Technol.* **4**, 014008 (2018).
- [26] J. M. Arrazola, O. Di Matteo, N. Quesada, S. Jahangiri, A. Delgado, and N. Killoran, Universal quantum circuits for quantum chemistry, *Quantum* **6**, 742 (2022).
- [27] G.-L. R. Anselmetti, D. Wierichs, C. Gogolin, and R. M. Parrish, Local, expressive, quantum-number-preserving VQE ansätze for fermionic systems, *New J. Phys.* **23**, 113010 (2021).
- [28] Y. S. Yordanov, D. R. M. Arvidsson-Shukur, and C. H. W. Barnes, Efficient quantum circuits for quantum computational chemistry, *Phys. Rev. A* **102**, 062612 (2020).
- [29] I. Magoulas and F. A. Evangelista, CNOT-efficient circuits for arbitrary rank many-body fermionic and qubit excitations, *J. Chem. Theory Comput.* **19**, 822 (2023).
- [30] C. H. Chee, A. M. Mak, D. Leykam, P. K. Barkoutsos, and D. G. Angelakis, Computing electronic correlation energies using linear depth quantum circuits, [arXiv:2207.03949](https://arxiv.org/abs/2207.03949).
- [31] I. Kerenidis and A. Prakash, Quantum machine learning with subspace states, [arXiv:2202.00054](https://arxiv.org/abs/2202.00054).
- [32] G. W. Stewart, Computing the CS decomposition of a partitioned orthonormal matrix, *Numer. Math.* **40**, 297 (1982).
- [33] E. S. Gawlik, Y. Nakatsukasa, and B. D. Sutton, A backward stable algorithm for computing the CS decomposition via the polar decomposition, *SIAM J. Matrix Anal. Appl.* **39**, 1448 (2018).
- [34] S. Johri, S. Debnath, A. Mocherla, A. Singk, A. Prakash, J. Kim, and I. Kerenidis, Nearest centroid classification on a trapped ion quantum computer, *npj Quantum Inf.* **7**, 1 (2021).
- [35] K. Wan, W. J. Huggins, J. Lee, and R. Babbush, Matchgate shadows for fermionic quantum simulation, [arXiv:2207.13723](https://arxiv.org/abs/2207.13723).
- [36] G. Ortiz, J. E. Gubernatis, E. Knill, and R. Laflamme, Quantum algorithms for fermionic simulations, *Phys. Rev. A* **64**, 022319 (2001).
- [37] Z. Jiang, K. J. Sung, K. Kechedzhi, V. N. Smelyanskiy, and S. Boixo, Quantum Algorithms to Simulate Many-Body Physics of Correlated Fermions, *Phys. Rev. Appl.* **9**, 044036 (2018).
- [38] A. Vourdas, Exterior calculus and fermionic quantum computation, *J. Phys. A* **51**, 445301 (2018).
- [39] I. Kerenidis, J. Landman, and N. Mathur, Classical and quantum algorithms for orthogonal neural networks, [arXiv:2106.07198](https://arxiv.org/abs/2106.07198).
- [40] J. D. Whitfield, J. Biamonte, and A. Aspuru-Guzik, Simulation of electronic structure Hamiltonians using quantum computers, *Mol. Phys.* **109**, 735 (2011).
- [41] M. Kjaergaard, M. E. Schwartz, J. Braumüller, P. Krantz, J. I.-J. Wang, S. Gustavsson, and W. D. Oliver, Superconducting qubits: Current state of play, *Annu. Rev. Condens. Matter Phys.* **11**, 369 (2020).
- [42] C. D. Bruzewicz, J. Chiaverini, R. McConnell, and J. M. Sage, Trapped-ion quantum computing: Progress and challenges, *Appl. Phys. Rev.* **6**, 021314 (2019).
- [43] J. M. L. Martin, Ab initio total atomization energies of small molecules — Towards the basis set limit, *Chem. Phys. Lett.* **259**, 669 (1996).
- [44] A. Halkier, T. Helgaker, P. Jørgensen, W. Klopper, and J. Olsen, Basis-set convergence of the energy in molecular Hartree–Fock calculations, *Chem. Phys. Lett.* **302**, 437 (1999).
- [45] P. R. Spackman and A. Karton, Estimating the CCSD basis-set limit energy from small basis sets: Basis-set extrapolations vs additivity schemes, *AIP Adv.* **5**, 057148 (2015).
- [46] C. Plascencia, J. Wang, and A. K. Wilson, Importance of the ligand basis set in ab initio thermochemical calculations of transition metal species, *Chem. Phys. Lett.* **685**, 496 (2017).
- [47] P. Virtanen *et al.*, SciPy 1.0: Fundamental algorithms for scientific computing in Python, *Nat. Methods* **17**, 261 (2020).
- [48] Q. Sun, T. C. Berkelbach, N. S. Blunt, G. H. Booth, S. Guo, Z. Li, J. Liu, J. D. McClain, E. R. Sayfutyarova, S. Sharma, S. Wouters, and G. K.-L. Chan, PySCF: The Python-based simulations of chemistry framework, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **8**, e1340 (2018).
- [49] V. Bergholm *et al.*, PennyLane: Automatic differentiation of hybrid quantum-classical computations, [arXiv:1811.04968](https://arxiv.org/abs/1811.04968).