

Facile synthesis of enol ethers *via* Zn(OTf)₂-mediated formal alkyne hydration-Smiles rearrangement†

Cite this: DOI: 10.1039/c4ra00159a

Received 8th January 2014
Accepted 25th March 2014

DOI: 10.1039/c4ra00159a

www.rsc.org/advances

Xinying Chew,^a Yuhan Lin^b and Yee Hwee Lim^{*a}

An efficient protocol involving commercially available zinc trifluoromethanesulfonate was able to transform a terminal alkyne to an enol ether *via* a formal tandem Markovnikov hydration-Smiles type rearrangement. The chemoselective reaction was able to work with a range of heteroaromatic tethers such as diazine, pyridine, benzimidazole and benzothiazole.

Alkyne hydration or the addition of water to an alkyne is an atom-economical¹ method to generate useful carbonyl compounds and therefore is a subject of much interest. Since the discovery of mercury(II) salts that hydrate alkynes with Markovnikov selectivity to give ketones,² most of the research has focused on the development of less toxic, more selective and efficient metal and acid catalysts or biocatalysts for this process.³ Non-metal based catalysts such as formic acid,⁴ triflic acid⁵ or trifluoroacetic acid⁶ can be used to hydrolyze terminal alkynes to the Markovnikov ketone product at elevated temperatures in water. Numerous metal salts and complexes have also been identified to convert terminal alkynes to ketones (Markovnikov product) or aldehydes (anti-Markovnikov product), (Scheme 1) with the most prominent being palladium(II),⁷ gold(I),^{7e,8} gold(III),⁹ platinum(II)¹⁰ and ruthenium¹¹ catalysts. Catalytic anti-Markovnikov alkyne hydration is a relatively new and exciting area where ruthenium¹² has emerged to be a highly efficient catalyst.


Being in the same group as mercury, studies using zinc for alkyne hydrations was reported as early as 1909 by Kutscheroff^{2c} but was found to require very harsh conditions.¹³ There were few reported research using zinc thereafter. In 1994, an anti-

Markovnikov hydration using silica gel-supported zinc borohydride to afford terminal alcohols rather than the aldehyde was disclosed.¹⁴


During the course of our investigations on diazines, we discovered an extremely facile protocol to convert the terminal alkynes to enol ethers using cheap and commercially available reagent, zinc triflate (Scheme 2). We believed that the terminal alkyne **1a** was first hydrated in a Markovnikov manner to give an intermediate **A** that underwent an intramolecular Smiles-type rearrangement¹⁵ to yield the diazine-protected enol ether **2a**. No ketone was ever observed or isolated.

An initial experiment was performed using **1a** with Zn(OTf)₂ in MeCN at ambient temperature. To our pleasant surprise, the reaction proceeded cleanly to give new product in nearly quantitative yield (95%) after 16 h. After extensive NMR, mass and IR spectroscopic analyses, the product was characterized to be **2a**.¹⁶ As Zn(OTf)₂ is not known to mediate alkyne hydration nor Smiles rearrangement to the best of our knowledge, we decided to further investigate this serendipitous observation.

A range of metal salts and acids commonly known to effect alkyne hydrations were first screened to determine if the



Scheme 1 General scheme of terminal alkyne hydration.



Scheme 2 Zinc-mediated formal alkyne hydration-Smiles rearrangement of diazine **1a**.

^aOrganic Chemistry, Institute of Chemical and Engineering Sciences (ICES), Agency for Science, Technology and Research (A*STAR), 11 Biopolis Way, #03-08 Helios Block, Singapore 138667, Singapore. E-mail: lim_yee_hwee@ices.a-star.edu.sg

^bSingapore Bioimaging Consortium (SBIC), Agency for Science, Technology and Research (A*STAR), 11 Biopolis Way, #01-02 Helios Block, Singapore 138667, Singapore

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ra00159a

Table 1 Initial screening for the alkyne hydration-Smiles rearrangement^a

Entry	Metal salt/acid	Temp. (°C)	Conv. ^b (%)	Yield ^b (%)
1	Zn(OTf) ₂	25	60	50
2	AgOTf	25	15	10
3	CuOTf	25	45	25
4	FeCl ₃	25	0	0
5	PdCl ₂	25	100	0
6	CuF ₂	25	10	0
7	RuCl ₃	25	100	0
8 ^c	TfOH	80	16	7
9	AgNTf ₂	80	80	40
10	AgOTf	80	50	45
11	CuOTf	80	100	40
12	Zn(OTf) ₂	80	100	95
13 ^d	Zn(OTf) ₂	80	48	45

^a Conditions: **1a** (1 equiv.), metal salt/acid (1 equiv.), MeCN (0.1 M), 2 h; then H₂O, 80 °C, 1 h. ^b Determined by ¹H NMR with 2,4,6-collidine as an internal standard. ^c 5 mol% of TfOH was used instead. ^d 50 mol% of Zn(OTf)₂ was used instead.

observed reactivity behavior was exclusive (Table 1).¹⁷ It was found that a variety of other metal salts such as AgNTf₂, AgOTf¹⁸ and CuOTf could yield the same product **2a** (Table 1, entries 2, 3

Table 2 Solvent optimization for the alkyne hydration-Smiles rearrangement^a

CC(C#CCOC1=CC=C(N=N)C=C1Cl) >> CC(C=CC(O)OC1=CC=C(N=N)C=C1Cl)

1a **2a**

Entry	Solvent	Conv. ^b (%)	Yield ^b (%)
1	MeCN	100	95
2	MeOH	15	10
3	DMSO	0	0
4	Acetone	100	25
5	1,4-Dioxane	100	20
6	EtOAc	100	40
7	H ₂ O	10	5
8	MeCN–H ₂ O (9 : 1)	100	75
9	MeCN–H ₂ O (4 : 1)	90	70
10	MeCN–H ₂ O (1 : 1)	75	55
11	MeCN–H ₂ O (1 : 4)	0	0
12	MeCN–H ₂ O (1 : 9)	0	0

^a Conditions: **1a** (1 equiv.), Zn(OTf)₂ (1 equiv.), solvent (0.1 M), 80 °C, 2 h; then H₂O, 80 °C, 1 h. ^b Determined by ¹H NMR with 2,4,6-collidine as an internal standard.

and 9–11). A control experiment with triflic acid was carried out to determine whether the reaction was mediated by Brønsted acid and only a small amount (7%) of the desired product could be obtained with catalytic amount of TfOH (Table 1, entry 8). Zn(OTf)₂ was by far superior in terms of efficiency and yield (Table 1, entries 1 and 12). At elevated temperature (80 °C), the reaction could be completed within 2 h to give **2a** in 95% NMR yield (Table 1, entry 12). Stoichiometric amount of Zn(OTf)₂ was necessary to effect the reaction as the conversion and yield dropped drastically with sub-stoichiometric amount of Zn(OTf)₂ (Table 1, entry 13).

Table 3 Hydration-rearrangement of diazines **1a**

Entry	Substrate	Product	T ^b (h)	Yield ^c (%)
1	1a	2a	2	97
2	1b	2b	2	95
3	1c	2c	2	95
4	1d	2d	3	88
5	1e	2e	3	86
6	1f	2f	3.5	85
7	1g	2g^d	3	87

^a Conditions: substrate **1** (1 equiv.), Zn(OTf)₂ (1 equiv.), MeCN (0.1 M), 80 °C, for specified time; then H₂O (1 mL mmol⁻¹), 80 °C for 1 h. ^b Reaction was monitored by TLC until all starting material was consumed. ^c Isolated yields. ^d The dehydrated product **2g** was isolated instead of the expected tertiary alcohol.

Different polar solvents were screened to determine the effect of solvents on the reaction (Table 2). Whilst solvents like MeOH, acetone, 1,4-dioxane and EtOAc were found to give some conversion (Table 2, entries 2, 4, 5 and 6), MeCN was found to be the best solvent. It is interesting to note that the reaction proceeded even in pure water, albeit with very low conversion (Table 2, entry 7). We then studied the water tolerance level of this reaction (Table 2, entries 8–12). For reactions in a mixture of MeCN and water, the efficiency of the reaction decreases as the water content increases. At 1 : 1 MeCN–H₂O mixture, the conversion dropped to 75% whilst the yield was only 55% (Table 2, entry 10).

In order to confirm that an external molecule of water was added during the hydration-rearrangement, **1c** was treated with Zn(OTf)₂ in MeCN at 80 °C for 2 h followed by a few drops of enriched ¹⁸O[H₂O]. Indeed, mass spectroscopic analysis of the product ¹⁸O-**2c** found incorporation of heavy water, affording a product of *m/z* 225.0278 for [M + Na]⁺.¹⁹ Based on the above

optimized conditions of using stoichiometric amount of Zn(OTf)₂ in MeCN at 80 °C, the scope of the reaction was then examined.²⁰

A range of diazines **1** were explored (Table 3). In general, the reactions proceeded smoothly with excellent yields (85–97%) to afford the corresponding enol ethers. The reaction was found to be tolerant of ether (Table 3, entry 4) as well as alkene functional groups (Table 3, entry 7). As expected, the stereochemistry of **1d** was retained in the hydration-rearrangement process (Table 3, entry 4).²¹ Excellent chemoselectivity was observed as demonstrated by substrate **1f** where the integrity of the second alkyne remained intact under the reaction conditions (Table 3, entry 6). The expected tertiary alcohol product for substrate **1g** was not obtained; instead the dehydrated triene **2g** was isolated (Table 3, entry 7).

The presence of the diazine was vital for the success of the hydration-rearrangement reaction. Apart from diazine as the tether for the hydration-rearrangement, we decided to explore if other similar aromatic groups could be used. Numerous functionalities were attempted, Table 4 detailed some of the results.

The pyridinyl **3a** proceeded smoothly to afford the corresponding enol ether **4a** in excellent yields (89%) (Table 4, entry 1). Benzoxazole ether **3b** did not react while the corresponding benzoxazole thioether **3c** gave the expected product **4c** in 45% yield together with another by-product **4c'** in 25% (Table 4, entries 2 and 3). **4c'** was presumed to be formed through a transfer of the benzoxazole group from another starting molecule **3c**. A similar reaction involving benzothiazole thioether **3d** afforded the expected enol ether **4d** in 92% yield in 6 h (Table 4, entry 4). Similarly, the benzimidazole thioether **3e** proceeded smoothly to give **4e** in 81% yield (Table 4, entry 5). No side products were observed in both cases. This difference in reactivity remains unclear at this point and is subject to further investigation.




Table 4 Other successful hydration-rearrangements^a

Entry	Substrate	Product	<i>T</i> ^b (h)	Yield ^c (%)
1			24	89
2			24	0
3			3	45 ^d
4			6	92
5			6	81

^a Conditions: substrate **3** (1 equiv.), Zn(OTf)₂ (1 equiv.), MeCN (0.1 M), 80 °C, for specified time; then H₂O (1 mL mmol⁻¹), 80 °C for 1 h.
^b Reaction was monitored by TLC until all starting material was consumed. ^c Isolated yields. ^d 25% of side product **4c'** was obtained.

Conclusions

In summary, a mild and efficient protocol to transform terminal alkynes to enol ethers *via* a formal tandem Markonikov hydration-Smiles rearrangement using cheap and commercially available zinc triflate has been developed. The reaction is highly chemoselective and works in a range of heteroaromatic such as diazine, pyridine, benzimidazole, benzoxazole and benzothiazole. The present study also highlights Zn(OTf)₂ as an eco-friendly reagent where the transformation is 100% atom-economical. Further investigations into the mechanism as well as other applications in synthesis are currently in progress.

Acknowledgements

We thank Ms Doris Tan (ICES) for high resolution mass spectrometric (HRMS) assistance. Financial support for this work

was provided by A*STAR Joint Council Organisation (JCO), Singapore under the JCO Career Development Award (CDA) grant JCO12302EG013 (to Y.H.L.). Dr Romain Bejot was gratefully acknowledged for helping to obtain a sample of enriched $^{18}\text{O}[\text{H}_2\text{O}]$ from Singapore Radiopharmaceuticals Ltd for our investigations.

Notes and references

- 1 B. M. Trost, *Science*, 1991, **254**, 1471.
- 2 (a) M. Kutscheroff, *Chem. Ber.*, 1881, **14**, 1540; (b) M. Kutscheroff, *Chem. Ber.*, 1884, **17**, 13; (c) M. G. Kutscheroff, *Chem. Ber.*, 1909, **42**, 2759.
- 3 L. Hintermann and A. Labonne, *Synthesis*, 2007, 1121.
- 4 (a) N. Menashe, D. Reshef and Y. Shvo, *J. Org. Chem.*, 1991, **56**, 2912; (b) N. Menashe and Y. Shvo, *J. Org. Chem.*, 1993, **58**, 7434.
- 5 T. Tsuchimoto, T. Joya, E. Shirakawa and Y. Kawakami, *Synlett*, 2000, 1777.
- 6 (a) F. Persico and B. D. Wuest, *J. Org. Chem.*, 1993, **53**, 95; (b) B. Meseguer, D. Alonso-Diaz, N. Griebenow, T. Herget and H. Waldmann, *Angew. Chem., Int. Ed.*, 1999, **38**, 2902; (c) S. Duclos, H. Stoeckli-Evans and T. R. Ward, *Helv. Chim. Acta*, 2001, **84**, 3148.
- 7 (a) A. Avshu, R. D. O'Sullivan, A. W. Parkins, N. W. Alcock and R. M. Countryman, *J. Chem. Soc., Dalton Trans.*, 1983, 1619; (b) K. Utimoto, *Pure Appl. Chem.*, 1983, **55**, 1845; (c) K. Imi, K. Imai and K. Utimoto, *Tetrahedron Lett.*, 1987, **28**, 3127; (d) Y. Kataoka, O. Matsumoto, M. Ohashi, T. Yamagata and K. Tani, *Chem. Lett.*, 1994, 1283; (e) B. Liu and J. K. De Brabander, *Org. Lett.*, 2006, **8**, 4907.
- 8 (a) E. Mizushima, K. Sato, T. Hayashi and M. Tanaka, *Angew. Chem., Int. Ed.*, 2002, **41**, 4563; (b) P. Roembke, H. Schmidbaur, S. Cronje and H. Raubenheimer, *J. Mol. Catal.*, 2004, **212**, 35; (c) E. Mizushima, D. Cui, D. C. D. Nath, T. Hayashi and M. Tanaka, *Org. Synth.*, 2006, **83**, 55.
- 9 (a) R. O. C. Norman, W. J. E. Parr and C. B. Thomas, *J. Chem. Soc., Perkin Trans. 1*, 1976, 1983; (b) Y. Fukuda and K. Utimoto, *J. Org. Chem.*, 1991, **56**, 3729; (c) R. Casado, M. Contel, M. Laguna, P. Romero and S. Sanz, *J. Am. Chem. Soc.*, 2003, **125**, 11925.
- 10 (a) J. Chatt, R. G. Guy and L. A. Duncanson, *J. Chem. Soc.*, 1961, 827; (b) W. Hiscox and P. W. Jennings, *Organometallics*, 1990, **9**, 1997; (c) J. W. Hartman, W. C. Hiscox and P. W. Jennings, *J. Org. Chem.*, 1993, **58**, 7613.
- 11 (a) J. Halpern, B. R. James and A. L. W. Kemp, *J. Am. Chem. Soc.*, 1961, **83**, 4097; (b) J. Halpern, B. R. James and A. L. W. Kemp, *J. Am. Chem. Soc.*, 1966, **88**, 5142; (c) Y. Sasaki and P. H. Dixneuf, *J. Chem. Soc., Chem. Commun.*, 1986, 790; (d) M. M. Taqui Khan, S. B. Halligudi and S. Shukla, *J. Mol. Catal.*, 1990, **58**, 299.
- 12 For selected references, see: (a) M. Tokunaga and Y. Wakatsuki, *Angew. Chem., Int. Ed.*, 1998, **37**, 2867; (b) M. Tokunaga and Y. Wakatsuki, Jap. Patent 11319576 A2, 1999, Chem. Abstr., 1999, 131, 352828; (c) M. Tokunaga, T. Suzuki, N. Koga, T. Fukushima, A. Horiuchi and Y. Wakatsuki, *J. Am. Chem. Soc.*, 2001, **123**, 11917; (d) D. B. Grotjahn, C. D. Incarvito and A. L. Rheingold, *Angew. Chem., Int. Ed.*, 2001, **40**, 3884; (e) D. B. Grotjahn and D. A. Lev, *J. Am. Chem. Soc.*, 2004, **126**, 12232; (f) F. Boeck, T. Kribber, L. Xiao and L. Hintermann, *J. Am. Chem. Soc.*, 2011, **133**, 8138.
- 13 M. Ban and F. Ida, *US Pat.*, 2791614, 1957, Chem. Abstr., 1957, 51, 85804.
- 14 B. C. Ranu, A. Sarkar, M. Saha and R. Chakraborty, *Tetrahedron*, 1994, **50**, 6579–6584.
- 15 (a) L. A. Warren and S. Smiles, *J. Chem. Soc., Abstr.*, 1930, 1327; (b) W. E. Truce, E. M. Kreider and W. W. Brand, *Org. React.*, 1970, **18**, 99.
- 16 For more details on the structural assignment of **2a**, please see ESI.†
- 17 For detailed screening results, please see ESI.†
- 18 R. Das and D. Chakraborty, *Appl. Organomet. Chem.*, 2012, **26**, 722.
- 19 **2c**: $\text{C}_8\text{H}_9\text{N}_2\text{O}_2\text{ClNa}$ has $[\text{M} + \text{Na}]^+$ calcd: 223.0245, found: 223.0249. ^{18}O -**2c**: $\text{C}_8\text{H}_9\text{N}_2\text{O}_{18}\text{ClNa}$ has $[\text{M} + \text{Na}]^+$ calcd: 225.0293, found: 225.0278.
- 20 Representative procedure: To a solution of **1a** (182 mg, 1 mmol) in MeCN (10 ml) was added $\text{Zn}(\text{OTf})_2$ (365 mg, 1 mmol) and the reaction mixture was stirred at 80 °C until consumption of **1a**. Water (1 mL) was then added and the reaction was stirred for another 1 h. Solvent was removed and the crude was diluted with EtOAc and washed with saturated NaHCO_3 . The crude was purified by column chromatography to afford the desired product **2a** as a colorless oil (95%).
- 21 See ESI† for more details on determination of absolute configuration.