

The glycaemic index and insulinaemic index of commercially available breakfast and snack foods in an Asian population

Wei Shuan Kimberly Tan¹, Wei Jie Kevin Tan¹, Shalini D/O Ponnalagu¹, Katie Koecher², Ravi Menon², Sze-Yen Tan¹ and Christiani J. Henry^{1,3*}

¹Clinical Nutrition Research Centre (CNRC), Singapore Institute for Clinical Sciences (SICS), Agency for Science, Technology and Research (A*STAR) and National University Health System, 117599 Singapore

²General Mills, Inc., Golden Valley, MN 55427, USA

³Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, 117597 Singapore

(Submitted 29 November 2017 – Final revision received 26 January 2018 – Accepted 23 February 2018)

Abstract

A low-glycaemic-index (GI) breakfast has been shown to lower blood glucose levels throughout the day. A wide variety of breakfast foods are consumed, but their GI values are largely unknown, hence limiting consumers' ability to select healthier options. This study investigated the GI values of ten common breakfast (five Asian and five Western) foods in this region using a randomised, cross-over study design. Participants arrived after an overnight fast, and fasting blood sample was taken before participants consumed test foods. Next, blood samples were taken at fixed intervals for 180 min. Glycaemic and insulinaemic responses to test foods were calculated as incremental AUC over 120 min, which were subsequently reported as glycaemic and insulinaemic indices. In all, nineteen healthy men (nine Chinese and ten Indians) aged 24·7 (SEM 0·4) years with a BMI of 21·7 (SEM 0·4) kg/m² completed the study. Asian breakfast foods were of medium (white bun filled with red bean paste = 58 (SEM 4); Chinese steamed white bun = 58 (SEM 3)) to high GI (rice idli = 85 (SEM 4); rice dosa = 76 (SEM 5); upma = 71 (SEM 6)), whereas Western breakfast foods were all of low GI (whole-grain biscuit = 54 (SEM 5); whole-grain biscuit filled with peanut butter = 44 (SEM 3); whole-grain oat muesli = 55 (SEM 4); whole-grain oat protein granola = 51 (SEM 4); whole-grain protein cereal = 49 (SEM 3)). The GI of test foods negatively correlated with protein ($r_s = -0.366$), fat ($r_s = -0.268$) and dietary fibre ($r_s = -0.422$) (all $P < 0.001$). GI values from this study contribute to the worldwide GI database, and may assist healthcare professionals in recommending low-GI breakfast to assist in lower daily glycaemia among Asians who are susceptible to type 2 diabetes mellitus.

Key words: Glycaemic index: Glucose: Insulin: Dietary proteins: Dietary fats: Dietary fibres

In the past decade, the prevalence of type 2 diabetes mellitus has been increasing steadily⁽¹⁾. Studies have shown that Asian populations, especially South Asians⁽²⁾, have an increased risk in the development of obesity, insulin resistance, type 2 diabetes mellitus and cardiovascular diseases compared with Caucasians^(3–5). The risks are further amplified by the traditional high-carbohydrate diets and lower physical activity levels observed among Asians^(6,7). In addition to increasing physical activity, it is also important to identify dietary interventions to improve insulin sensitivity for better glucose management and prevention of diabetes. Consuming a low-glycaemic-index (GI) diet has been implicated in minimising the risk of developing type 2 diabetes mellitus^(8,9). A simple replacement of white rice (higher GI) with brown rice (lower GI) has shown to reduce 24-h glucose and fasting insulin responses in overweight Indian adults⁽¹⁰⁾. Similarly, the consumption of low-GI rice at breakfast showed a reduction in post-prandial glycaemic response (GR) and food intake during the subsequent meals⁽¹¹⁾.

GI is a method of characterising carbohydrate foods on the basis of how much these foods increase blood glucose levels relative to an equal amount of glucose over 2 h post ingestion⁽¹²⁾. Carbohydrate foods are classified as low (≤ 55), medium (56–69) and high (≥ 70) GI⁽⁹⁾. Higher-GI foods may influence glucose intolerance and/or development of type 2 diabetes. One of the proposed mechanisms is that the large incremental glucose concentration may exaggerate the body's normal anabolic responses⁽¹³⁾. This consequently facilitates overproduction of insulin, and over an extended period of repeated exposure it results in pancreatic β -cell deficiency⁽¹⁴⁾. In our previous studies on healthy Chinese male population, we have shown that consuming a high-GI breakfast increased blood glucose concentrations not only after breakfast, but also throughout the entire day compared with a matched low-GI breakfast. Moreover, it has been reported that low-GI breakfast foods resulted in a significantly lower GR, which had a second meal effect on lunch, and resulted in a significantly decreased

Abbreviations: GI, glycaemic index; GR, glycaemic response; iAUC, incremental AUC; II, insulinaemic index.

* **Corresponding author:** C. J. Henry, fax +65 6776 6840, email jeya_henry@sics.a-star.edu.sg

energy intake⁽¹¹⁾. This highlights the importance of consuming low-GI breakfast foods, which results in a day-long lower glycaemia. The insulinaemic index (II), similar to GI, is the insulin response to a carbohydrate-rich food. It is calculated as the incremental AUC (iAUC) of insulin from test food over 120 min compared with the iAUC of insulin from the glucose reference⁽¹⁵⁾. Excessive post-prandial insulin secretion has been shown to increase oxidative stress and accelerate the decline of β -cell function especially in individuals who are at risk of developing type 2 diabetes^(16,17). Hence, the measurement of II along with GI of food is an important biomarker of the metabolic consequences of consuming a carbohydrate-rich diet⁽¹⁷⁾. Although post-prandial glucose response is closely associated with insulin excursions, this association is not consistent especially in the presence of protein, fat, fibre and structural differences^(18–20).

As Singapore imports over 90% of its food owing to the limited farming grounds⁽²¹⁾, various traditional and international breakfast foods are easily and widely consumed in Singapore. Given the wide variety of breakfast cereals consumed in Singapore and this region, the GI of these breakfast foods are largely unknown. To provide scientific evidence on breakfast choices to the local populations, this study investigated the GI and II of selected Western and Asian breakfast foods consumed locally. As Western and Asian breakfast foods included in this study differed in macronutrient composition, this study also investigated how these nutrients may be associated and influence both glycaemic and insulinaemic response (IR).

Methods

Participants

The inclusion criteria were healthy Chinese and Indian men aged between 21 and 40 years, who were non-smokers with a healthy BMI between 18.5 and 25 kg/m². Participants with metabolic diseases (such as diabetes, hypertension and so on), known glucose-6-phosphate dehydrogenase deficiency, medical conditions and/or taking medications known to affect glycaemia (glucocorticoids, thyroid hormones, thiazide diuretics), intolerances or allergies to foods, those who partake in sports at the competitive and/or endurance levels, those who intentionally restrict food intake and those whose fasting blood glucose was more than 6 mmol/l and HbA1c was more than 6% were excluded from the study. A total of twenty-four

participants were screened and twenty-two participants (eleven Chinese and eleven Indians) were recruited. Two participants failed screening owing to BMI not being within range. The study was conducted in accordance with the guidelines laid down in the Declaration of Helsinki, and all procedures involving human participants were approved by the Domain Specific Review Board of National Healthcare Group, Singapore (reference no. 2016/00614). Participants gave their informed consent before their participation.

Study protocol

The study was conducted using a randomised, cross-over design. Participants were given the option to choose between five or ten breakfast test foods. For all participants, the first three test sessions were the reference foods (50 g of glucose anhydrous powder dissolved in 250 ml of water) followed by five or ten sessions of different breakfast test foods containing 50 g of available carbohydrate in randomised order (using <https://www.randomizer.org/>). Five Asian and five Western breakfast test foods were selected. The five Asian breakfast foods were commercially available dry mix rice idli (India), upma (India), rice dosa (India), white bun filled with red bean paste (Singapore) and Chinese steamed white bun (Malaysia), whereas the five Western breakfast were whole-grain biscuits (USA), whole-grain biscuit filled with peanut butter (USA), whole-grain oat muesli (USA), whole-grain oat protein granola (USA) and whole-grain protein cereal (USA). These foods were chosen for their variety, cultural relevance (i.e. snacks or breakfast) and were commonly consumed in Asia. Moreover, their GI values were not reported in any established GI database. The foods were selected to represent a wide spectrum of grain type, macronutrient composition and differing form (flaked oats in muesli, milled rice in idli and so on). These are some of the factors that influence GI and II.

Nutrient analyses were conducted for all test foods, including available carbohydrate content assessed by Medallion Laboratories, and were served in portions containing 50 g of available carbohydrates (Table 1). The food was prepared freshly as per cooking instructions given on the packaging (Table 2) every morning and was served with 250 ml of water. All foods tested had $\geq 45\%$ of energy content coming from carbohydrates.

Participants arrived at the centre at 08.30 hours after a 10- to 12-h overnight fast. They were advised to follow a standard

Table 1. Serving portion sizes and nutrient compositions of ten test foods (per serving of 50 g of available carbohydrate product)

Test foods	Serving size (g)	Energy (kJ)	Fat (g)	Protein (g)	Dietary fibre (g)
Rice idli	162	1075	2.7	6.4	3.9
Upma	310	1276	7.8	6.9	5.1
Rice dosa	193	1477	12.8	7.4	5.5
White bun filled with red bean paste	106	1469	12.9	6.5	5.2
Chinese steamed white bun	88	1046	2.6	5.8	4.4
Whole-grain biscuits	82	1602	14.8	7.5	9.9
Whole-grain biscuit filled with peanut butter	102	2180	26.9	12.8	11.9
Whole-grain oat muesli	84	1523	11.8	9.8	9.0
Whole-grain oat protein granola	87	1561	9.4	18.3	7.1
Whole-grain protein cereal	99	1711	10.4	28.8	5.6

Table 2. Preparation method and main ingredients of ten commercially available Western and Asian breakfast foods

Foods	Preparation method for 50 g of available carbohydrate	Main ingredients
Rice idli	Mix 70.0 g of idli premix to 91.0 g of water. Grease the idli mould with 0.75 g of ghee and steam for 15 min	Rice flour, urad dal (dehusked split black lentil) flour and baking soda
Upma	Boil 235.0 g of water and stir in 75.2 g of upma premix. Stir constantly and serve after 3 min	Semolina, vegetable oil, salt, urad dal (dehusked split black lentil) and green chillies
Rice dosa	Mix 72.8 g of dosa premix and 110.0 g of water until smooth. Pour batter on a preheated pan and add 15 g of melted ghee (clarified butter). Flip the dosa after 1 min and cook for another minute. Serve	Rice flour, wheat flour, urad dal (dehusked split black lentil) flour, palm oil and salt
White bun filled with red bean paste	Microwave buns for 1 min and weigh 105.7 g of white bun filled with red bean paste	Wheat flour, sugar, red bean, palm olein and yeast
Chinese steamed white bun	Microwave buns for 1 min and weigh 87.9 g of white bun	Wheat flour, water, sugar, palm vegetable fat and baking powder
Whole-grain biscuits	Weigh 83.6 g of biscuit and serve	Whole-grain oats, rapeseed oil, sugar, whole-grain barley flakes and honey
Whole-grain biscuit filled with peanut butter	Weigh 101.8 g of biscuit and serve	Whole-grain oats, sugar, peanut butter (peanuts, salt), whole-grain barley flakes and rapeseed oil
Whole-grain oat muesli	Weigh 83.6 g of muesli and serve	Whole-grain oats, sugar, sunflower seeds, dried cranberries and raisins
Whole-grain oat protein granola	Weigh 87.3 g of granola and serve	Whole-grain oats, sugar, soya protein isolate, honey and rice starch
Whole-grain protein cereal	Weigh 56.9 g of cereal and serve	Whole-grain oats, soya protein isolate, whole-grain oats clusters, sugar and almonds

dinner, and to avoid any strenuous physical activities and consumption of alcoholic beverages the day before the test sessions. After a 10-min rest, two finger-prick blood samples were collected 5 min apart to measure baseline blood glucose concentrations, and approximately 0.5 ml of blood sample was collected for insulin analysis. Another fasting blood sample was taken if the first two baseline readings were more than 0.2 mmol/l apart. Subsequently, participants were given either a reference glucose drink or a test food to be consumed within 12 min. After consumption of breakfast test foods, approximately 5 µl of capillary blood samples were collected at 15-, 30-, 45-, 60-, 90-, 120-, 150- and 180-min time points for glucose measurements, and 0.5 ml of blood sample was collected at similar time points for insulin measurements.

The methodology used to measure the GI was adopted from that described by Brouns *et al.*⁽²²⁾ and was in line with procedures recommended by the ISO 26642⁽²³⁾. The blood glucose concentration in the sample was measured and recorded using the HemoCue® 201+ Glucose analyser (HemoCue Ltd)⁽²⁴⁾. The mean intra- and inter-assay CV for glucose were 1.2 and 1.3%, respectively.

For the measurement of blood insulin levels at baseline and every 30-min intervals, 500 µl of capillary blood (obtained from fingerprick) was collected in a Microtainer containing EDTA and chilled. Fingerpricks were made using the Accu-Check single-use lancing device (Roche). To minimise plasma dilution, fingertips were gently massaged starting from the base of the hand moving towards the tips. The blood samples were centrifuged at 6000 g for 10 min and the supernatant plasma was pipetted into Eppendorf tubes and stored in a -80°C freezer for further analysis. Insulin concentration (µU/ml) in the plasma samples was determined using the Cobas e411 immunochemistry analyzer (Roche Diagnostics). The mean intra- and inter-assay CV for insulin were 1.9 and 2.6%, respectively.

Statistical analysis

To obtain sufficient statistical power for GI testing of a single food, a minimum of ten participants are recommended by the ISO26642⁽²³⁾ standard and by Brouns *et al.*⁽²²⁾. A total of nineteen participants were therefore used in the data analysis. With reference to the present study, retrospective power calculation for a repeated measures design with ten treatments and nineteen participants, to detect significant difference between treatments with partial η^2 of 0.40 and non-sphericity correction, ϵ , of 0.6 resulted in a power larger than 95%. Moreover, using the power calculation with 80% power at 0.05 significant level, a minimum sample size of six participants would be required in the future. This indicates that sufficient participants were recruited for this present study.

The increment in GR was calculated by taking the difference of blood glucose reading at each time point and mean baseline value. The total GR over 120 min was also calculated as iAUC, using the trapezoidal rule that ignored the area under the baseline⁽²⁵⁾. In the case of the reference food, the mean of the three glucose tests was taken as the iAUC for the reference food. If the mean within-participant CV for three glucose drinks was >30%, outlying values were excluded as per standard protocol⁽²³⁾. Using the iAUC values, the GI of the test food was calculated using the following formula⁽²⁶⁾:

$$GI = \left(\frac{\text{iAUC of plasma glucose for the test food} / \text{average iAUC of}}{\text{plasma glucose for the glucose reference} \times 100} \right)$$

The calculation of II is similar to that of GI, that is II = (iAUC of plasma insulin for the test food/average iAUC of plasma insulin for the glucose reference × 100).

The final GI value of the test foods was determined by taking the average GI values of a test food from all participants. Participants whose GI value was not within two standard

Table 3. Anthropometric characteristics of the study participants (*n* 19, nine Chinese, ten Indians) (Mean values with their standard errors)

Anthropometric and physiological parameters	Mean	SEM
Age (years)	24.7	0.4
Height (cm)	173.7	1.0
Weight (kg)	65.6	1.5
BMI (kg/m ²)	21.7	0.4
Waist circumference (cm)	77.5	1.3
Hip circumference (cm)	95.7	1.2
Fasting blood glucose (mmol/l)	4.3	0.1
HbA1c (%)	5.3	0.1

deviations from the mean GI value were excluded as per standard GI protocol⁽²³⁾. The same procedures were repeated for IR and II. Simple Spearman's correlation and partial correlation controlling for two macronutrient contents at each time was also performed to evaluate the strength of association between the GI values and macronutrient content of the test foods. All statistical analyses were performed using SPSS version 24.0 (IBM Corp.), and statistical significance was set at $P < 0.05$, two-tailed.

Results

A total of twenty-two participants were recruited (eleven Chinese and eleven Indians). Three participants were excluded from the analysis owing to incomplete test sessions (one Chinese), withdrawal of consent owing to time commitment issues (one Indian) and GR that was deemed outliers based on the ISO 26642 definition⁽²³⁾. The final analysis of this study therefore included nineteen participants, and the anthropometric characteristics of these participants were reported in Table 3. The mean within-participant CV of GR and IR (based on iAUC) to glucose standard drinks for nineteen participants were 14.5 and 15.4%, respectively.

Temporal changes in capillary glucose after the consumption of test meals were presented in Fig. 1. The GI of ten test foods calculated on the basis of the iAUC are presented in Table 4. Rice idli, upma and rice dosa were categorised as high-GI foods; Chinese steamed white bun and white bun filled with red bean were categorised as medium-GI foods; and whole-grain biscuit, whole-grain biscuit filled with peanut butter, whole-grain oat muesli, whole-grain oat granola and whole-grain protein cereal were categorised as low-GI foods. The GI values of test breakfast foods were significantly and negatively correlated with overall protein ($r_s = -0.366$, $P < 0.001$), fat ($r_s = -0.268$, $P < 0.001$) and dietary fibre ($r_s = -0.422$, $P < 0.001$) levels. When partial correlations were performed controlling for two other macronutrients each time, only dietary fibre content was observed to maintain its negative correlation with GI (partial correlation between GI and dietary fibre controlling for protein and fat) ($r_s = -0.218$, $P = 0.003$). As for insulin, the post-prandial changes in insulin concentrations after meals are shown in Fig. 2 and the II of test foods are reported in Table 4. There was a significant negative correlation observed between II and overall dietary fibre ($r_s = -0.168$, $P = 0.021$), but not with fat and protein content.




Fig. 1. (a) and (b) Temporal curves of blood glucose response to different breakfast foods containing 50 g of available carbohydrates. Values are means with their standard errors. a: \blacklozenge , Glucose; \square , rice idli; \cdots , upma; \times , rice dosa; \triangle , white bun filled with red bean paste; \bullet , Chinese steamed white bun; b: \blacklozenge , glucose; \blacksquare , whole-grain biscuit; \blacktriangle , whole-grain biscuit filled with peanut butter; \cdots , whole-grain oat muesli; $----$, whole-grain oat protein granola; \ominus , whole-grain protein cereal.

Discussion

We tested the GI and II of ten commercially available Western and Asian breakfast foods in both Chinese and Indian participants. Among the ten test foods, five were of low GI, two were of medium GI and three were of high GI. Interestingly, out of all the test foods, all medium- and high-GI breakfast foods were of Asian cuisines, whereas all low-GI breakfast foods were of Western cuisines.

The high GI of both rice dosa and rice idli may be partly attributed to the larger proportion of refined rice flour compared with de-husked split black lentil flour in these foods. Refined rice has been reported to increase GI⁽²⁷⁾. This was evident from the steep glucose response peak observed at 30 min and high GI classification (Fig. 1(a)). In addition, these foods had low fat and protein content leading to a rapid increase in GR. Significant negative correlations found between GI and fat, protein and dietary fibre content supported this speculation. Moreover, our previous study has reported significant reduction in the GI of white rice when protein from chicken was added to the white rice⁽²⁸⁾. Studies have shown that the presence of fat significantly reduces GR by slowing gastric emptying⁽¹⁹⁾ and the presence of protein increases insulin secretion⁽²⁸⁾. This may explain why whole-grain protein cereal, whole-grain biscuit filled with peanut butter and whole-grain protein granola have low GI. However, for

whole-grain biscuit filled with peanut butter, it is postulated that the presence of higher fat content delayed gastric emptying. This reduces the rate of glucose uptake into the circulation, which

resulted in a reduced insulin response. Upma is an Indian dish that mainly consists of semolina, which is derived from wheat. Its consistency is similar to couscous. The GI of upma (GI = 71) was comparable to the GI of couscous (GI = 65 ± 4) that has previously been reported⁽¹²⁾.

Despite having low fat and protein content, Chinese steamed white bun was categorised as medium GI. This may be owing to the use of frozen Chinese white bun, as studies have shown that freezing may cause starch retrogradation and the formation of resistance starch^(29,30). This reduces starch digestion, which may reduce GR. This may also explain the lower GI value of white bun filled with red bean paste (thawed from frozen) compared with rice dosa while having similar protein and fat composition. Of the ten foods tested, it was found that three of the breakfast/snacks were of high GI, two were of medium GI and five were of low GI. In recommending the optimal breakfast choice, consumers are encouraged to select these five low-GI foods in the first instance. If medium- or high-GI foods are selected, a recommendation to reduce the portion size by approximately 20–30% will reduce the glycaemic load correspondingly⁽³¹⁾.

Whole-grain biscuit, whole-grain biscuit filled with peanut butter and whole-grain oat muesli had the highest dietary fibre content among all ten foods (Table 1). Certain dietary fibres play a role in the modulation of GR, which lowers the overall GI of those foods⁽³²⁾. The lower GI of whole-grain oat protein granola and muesli may also be explained by the structure of the carbohydrate. These products contained intact whole grains, which may lower the digestion and availability of carbohydrates. Studies have shown that the degree of processing or refining of grains results in structural differences that can have an impact on GR^(20,27). These examples showed that structural variations do affect the overall GI of the product in addition to nutrient composition.

In conclusion, this study provides GI and II values of ten commercially available Western and Asian breakfast foods. The presence of macronutrients, dietary fibre and level of grain intactness had significant influence in the overall GI of the breakfast foods. Between GI and II, it appears that II values markedly differ




Fig. 2. (a) and (b) Temporal curves of blood insulin response to different breakfast foods containing 50 g of available carbohydrates. Values are means with their standard errors. a: —◆—, Glucose; —□—, rice idli; upma; —×—, rice dosa; —△—, white bun filled with red bean paste; —●—, Chinese steamed white bun; b: —◆—, glucose; —■—, whole-grain biscuit; —▲—, whole-grain biscuit filled with peanut butter; —○—, whole-grain oat muesli; —○—, whole-grain oat protein granola; —○—, whole-grain protein cereal.

Table 4. Glycaemic (GI) and insulinaemic index (II) of ten breakfast foods (n 19) (Mean values with their standard errors)

Foods	GI		GI classification*	II	
	Mean	SEM		Mean	SEM
Rice idli†	85 ^a	4	High	114 ^{a,b,d}	10
Upma†	71 ^{a,b,c}	6	High	114 ^{b,d}	10
Rice dosa†	76 ^{a,b}	5	High	114 ^{a,b,d}	10
White bun filled with red bean paste‡	58 ^{b,d}	4	Med	98 ^{a,b,c}	11
Chinese steamed white bun†	58 ^{b,c,d}	3	Med	76 ^c	7
Whole-grain biscuits‡	54 ^{c,d}	5	Low	86 ^{c,d}	8
Whole-grain biscuit filled with peanut butter‡	44 ^d	3	Low	74 ^c	7
Whole-grain oat muesli‡	55 ^{b,d}	4	Low	83 ^{a,c}	8
Whole-grain oat protein granola‡	51 ^{c,d}	4	Low	111 ^{a,b,c}	12
Whole-grain protein cereal‡	49 ^d	3	Low	122 ^{a,b}	10

^{a,b,c,d}Foods that share the superscript letters are not significantly different from each other in terms of mean GI (mean II). However, the superscript letters in GI are not the same as the superscript letters in II. Repeated measures ANOVA was used to test for significant differences in mean GI (mean II) between foods.

* GI classification: low ≤ 55, medium: 56–69, high ≥ 70.

† Asian breakfast.

‡ Western breakfast.

between subjects, making II less universally valid than GI⁽²⁵⁾. Consumers are therefore best advised to consume foods with low GI. The results obtained may contribute to the evolving global database of GI of foods, and may assist in the selection of lower-GI breakfast/snacks to support day-long glycaemic control among Asians who are at increased risks for diabetes.

Acknowledgements

The authors thank the participants for their participation in this study.

This research was co-funded by General Mills Inc. and Singapore Institute for Clinical Sciences, A* STAR.

C. J. H., K. K. and R. M. designed the study and critically reviewed the manuscript. W. S. K. T. and W. J. K. T. assisted in study design, obtained study data, performed the data analysis and drafted the manuscript. S.-Y. T. and S. P. analysed the data and reviewed the manuscript. All authors approved the final manuscript as submitted.

C. J. H. is the guarantor of this work and had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. None of the authors has any conflicts of interest to declare.

References

- Roglic G (2016) WHO Global report on diabetes: a summary. *Int J Noncommun Dis* **1**, 3–8.
- Misra A & Khurana L (2011) Obesity-related non-communicable diseases: South Asians vs White Caucasians. *Int J Obes (Lond)* **35**, 167–187.
- Ramachandran A, Snehalatha C, Shetty AS, *et al.* (2012) Trends in prevalence of diabetes in Asian countries. *World J Diabetes* **3**, 110–117.
- Dickinson S, Colagiuri S, Faramus E, *et al.* (2002) Postprandial hyperglycemia and insulin sensitivity differ among lean young adults of different ethnicities. *J Nutr* **132**, 2574–2579.
- Chan JC, Malik V, Jia W, *et al.* (2009) Diabetes in Asia: epidemiology, risk factors, and pathophysiology. *JAMA* **301**, 2129–2140.
- Misra A, Khurana L, Isharwal S, *et al.* (2008) South Asian diets and insulin resistance. *Br J Nutr* **101**, 465–473.
- Phan TP, Alkema L, Tai ES, *et al.* (2014) Forecasting the burden of type 2 diabetes in Singapore using a demographic epidemiological model of Singapore. *BMJ Open Diabetes Res Care* **2**, e000012.
- Schulze MB, Liu S, Rimm EB, *et al.* (2004) Glycemic index, glycemic load, and dietary fiber intake and incidence of type 2 diabetes in younger and middle-aged women. *Am J Clin Nutr* **80**, 348–356.
- Willett W, Manson J & Liu S (2002) Glycemic index, glycemic load, and risk of type 2 diabetes. *Am J Clin Nutr* **76**, 274S–280S.
- Mohan V, Spiegelman D, Sudha V, *et al.* (2014) Effect of brown rice, white rice, and brown rice with legumes on blood glucose and insulin responses in overweight Asian Indians: a randomized controlled trial. *Diabetes Technol Ther* **16**, 317–325.
- Kaur B, Ranawana V, Teh A-L, *et al.* (2015) The impact of a low glycemic index (GI) breakfast and snack on daily blood glucose profiles and food intake in young Chinese adult males. *J Clin Transl Endocrinol* **2**, 92–98.
- Atkinson FS, Foster-Powell K & Brand-Miller JC (2008) International tables of glycemic index and glycemic load values: 2008. *Diabetes Care* **31**, 2281–2283.
- Ludwig DS (2002) The glycemic index: physiological mechanisms relating to obesity, diabetes, and cardiovascular disease. *JAMA* **287**, 2414–2423.
- Ceriello A (2005) Postprandial hyperglycemia and diabetes complications: is it time to treat? *Diabetes* **54**, 1–7.
- Wolever TMS, Jenkins DJA, Collier GR, *et al.* (1988) Metabolic response to test meals containing different carbohydrate foods: 1. Relationship between rate of digestion and plasma insulin response. *Nutr Res* **8**, 573–581.
- Nimptsch K, Brand-Miller JC, Franz M, *et al.* (2011) Dietary insulin index and insulin load in relation to biomarkers of glycemic control, plasma lipids, and inflammation markers. *Am J Clin Nutr* **94**, 182–190.
- Mimiran P, Esfandiari S, Bahadoran Z, *et al.* (2016) Dietary insulin load and insulin index are associated with the risk of insulin resistance: a prospective approach in Tehran Lipid and Glucose Study. *J Diabetes Metab Disord* **15**, 23.
- Alminger M & Eklund-Jonsson C (2008) Whole-grain cereal products based on a high-fibre barley or oat genotype lower post-prandial glucose and insulin responses in healthy humans. *Eur J Nutr* **47**, 294–300.
- Gentilcore D, Chaikomin R, Jones KL, *et al.* (2006) Effects of fat on gastric emptying of and the glycemic, insulin, and incretin responses to a carbohydrate meal in type 2 diabetes. *J Clin Endocrinol Metab* **91**, 2062–2067.
- Juntunen KS, Laaksonen DE, Autio K, *et al.* (2003) Structural differences between rye and wheat breads but not total fiber content may explain the lower postprandial insulin response to rye bread. *Am J Clin Nutr* **78**, 957–964.
- Agri-Food & Veterinary Authority of Singapore (2015) The food we eat. Singapore's Food Supply. <http://www.ava.gov.sg/explore-by-sections/food/singapore-food-supply/the-food-we-eat> (accessed August 2017).
- Brouns F, Bjorck I, Frayn KN, *et al.* (2005) Glycaemic index methodology. *Nutr Res Rev* **18**, 145–171.
- International Organization for Standardization (2010) *Food Products – Determination of the Glycaemic Index (GI) and Recommendation for Food Classification*. Geneva: International Organization for Standardization.
- Tan VM, Wu T, Henry CJ, *et al.* (2015) Glycaemic and insulin responses, glycaemic index and insulinaemic index values of rice between three Asian ethnic groups. *Br J Nutr* **113**, 1228–1236.
- Wolever TMS (2006) *The Glycaemic Index – A Physiological Classification of Dietary Carbohydrate*. Wallingford: CABI.
- Food and Agriculture Organization (1998) *Carbohydrates in Human Nutrition: Report of a Joint FAO/WHO Expert Consultation*. Rome: FAO.
- Heaton KW, Marcus SN, Emmett PM, *et al.* (1988) Particle size of wheat, maize, and oat test meals: effects on plasma glucose and insulin responses and on the rate of starch digestion *in vitro*. *Am J Clin Nutr* **47**, 675–682.
- Sun L, Ranawana DV, Leow MK, *et al.* (2014) Effect of chicken, fat and vegetable on glycaemia and insulinaemia to a white rice-based meal in healthy adults. *Eur J Nutr* **53**, 1719–1726.
- Burton P & Lightowler HJ (2007) The impact of freezing and toasting on the glycaemic response of white bread. *Eur J Clin Nutr* **62**, 594–599.
- Carreira MC, Lajolo FM & Menezes EWd (2004) Glycemic index: effect of food storage under low temperature. *Braz Arch Biol Technol* **47**, 569–574.
- Brand-Miller JC, Thomas M, Swan V, *et al.* (2003) Physiological validation of the concept of glycemic load in lean young adults. *J Nutr* **133**, 2728–2732.
- Russell WR, Baka A, Björck I, *et al.* (2016) Impact of diet composition on blood glucose regulation. *Crit Rev Food Sci Nutr* **56**, 541–590.